

Selenium levels in rice samples from high and low risk areas for esophageal cancer

Hadi Rahimzadeh-Barzoki, MSc,
Hamidreza Joshaghani, PhD, Somayeh Beirami, MSc,
Morteza Mansurian, MSc, Shabryar Semnani, MD,
Gholamreza Rosbandel, MD.

ABSTRACT

Objectives: To assess the relationship between selenium (Se) concentration in rice and the incidence of esophageal cancer (EC) in a high risk area in Northern Iran.

Methods: This ecological study was conducted in Golestan province of Iran in 2012. In this area, 45714 acres of land are cultivated by rice. A total of 69 rice samples were taken. We investigated Se concentrations by the voltammetric method. Statistical analysis was performed using the Pearson correlation test and Mann-Whitney U test.

Results: The mean (\pm SD) Se level in rice samples was 0.229 (\pm 0.145) mg/kg. The Se concentration was significantly higher in rice samples from high EC rate areas (0.35 mg/kg) compared with low risk areas (0.16 mg/kg) ($p < 0.001$). There was a significant positive correlation between the levels of Se in rice and the incidence rate of EC ($p = 0.03$).

Conclusion: We found a high rice Se concentration and a significant positive relationship between rice Se levels and EC rates in the Golestan province of Iran. High soil and rice Se levels may play a possible role in the pathogenesis of EC in this area.

Esophageal cancer (EC) is among eighth most common cancer in the world causing approximately 400,000 deaths in 2012.¹ Golestan province, in northeast Iran has been known as a high risk area for EC, most commonly esophageal squamous cell carcinoma (ESCC).² Esophageal cancer is a multi-factorial condition and various risk factors were proposed to play a role in its pathogenesis including high levels of selenium (Se) in serum³ and soil⁴ samples. Selenium is a metalloid and may be present in nature and in organisms as organic and/or inorganic forms. The essentiality and toxicity of Se are well known. Insufficient intake has been linked to serious health effects, such as dilated cardiomyopathy in Keshan disease, and osteoarthritis in Kashin-Beck disease.⁵ However, high exposure to Se may be toxic to human

beings.⁶ The most important source of Se for animals and humans is food, especially cereals.⁷ Rice is one of the commonly used cereals throughout the world. According to the Food and Agriculture Organization report,⁸ almost 30% of energy, and 20% of the world's protein supply is provided via the use of rice. Rice is the main food in Asian countries including Iran. Golestan Province of Iran is considered an important area in rice cultivation. In view of the high usage of rice and the importance of EC in the Golestan province of Iran, we conducted this study to assess any relationship between rice Se concentration and the incidence of EC in this area.

Methods. Collection of samples. This ecological study was conducted in Golestan province in northern Iran between April 2012 and August 2012. The area under rice cultivation is 45714 acres. Rice is a major crop in this region. A total of 69 rice samples were collected from different regions of Golestan province (42 samples from low EC-risk areas and 27 samples from high EC-risk areas, based on rice cultivation in each area) (Figure 1). Samples were collected from rice



Figure 1 - Map of the Golestan province indicating the number of rice samples collected from different regions and cities. Reprinted from Semnani S, Rosbandel G, Zendeabad A, Keshkar A, Rahimzadeh H, Abdolahi N, et al. Soils selenium level and esophageal cancer: an ecological study in a high risk area for esophageal cancer. *J Trace Elem Med Biol* 2010; 24: 174-177, with permission from Elsevier.

Disclosure. Authors have no conflict of interests, and the work was not supported or funded by any drug company. This article was part of a research project that was approved and supported by the Golestan University of Medical Sciences, Gorgan, Iran [project number: 9010130209].

farms, put in impermeable plastic bags and kept in a refrigerator until analyzing.

Reagents. All reagents were of analytical grade. Deionized water was used in preparing stock solutions. All glass laboratory equipment were previously soaked in 30% nitric acid (HNO_3) (v/v) for 24 hours and rinsed with distilled water.

Sample preparation. The rice samples were washed with deionized water, and then were dried by oven at 60°C for 24 hours. The dried samples were milled to fine powder. From the milled rice, 1g was weighed and digested in a high-walled beaker with a mixture solution of HNO_3 - H_2O_2 . A sample of rice (1 g) and 5 mL HNO_3 (65%) were transferred into a 100 mL beaker and kept overnight. It was evaporated gently by using a hot plate until its volume reached approximately one-third of the original volume. Heating was continued until the evolution of brown fumes of nitrogen oxides ceased. Then, 2 mLs of H_2O_2 were added and heated again. This solution was cooled to room temperature and filtrated through a Whatman filter (0.45 μ), (GE health care, Little Chalfont, UK) into a 100 mL volumetric flask and filled to 100 mL with deionized water. Solutions were transferred to a polyethylene terephthalate bottle and were kept in a refrigerator until Se was measured using a polarograph, (Metrohm AG, Herisau, Switzerland).

Selenium determination by the voltammetric method. The voltammetric analysis⁹ was carried out with a Metrohm 797 VA Computrace (Herisau, Switzerland). The 3 electrode classical cell consisted of an Hg electrode, a Pt electrode, and an Ag/AgCl reference electrode as well as a hanging mercury drop electrode (HMDE). Only high purity grade (Merck and Chem lab Suprapur, Darmstadt, Germany) reagents and deionized water were used. For calibration, the standard addition method⁹ was applied. The calibration standards were prepared using Chem Lab, Zedelgem, Belgium at 1000 ppm of Se. The Se concentrations were quantified using linear regression based on the height of the peaks of the voltammograms.

Data were entered into a computer and statistical analysis was carried out using the Statistical Package for Social Sciences, version 16 (SPSS Inc., Chicago, IL, USA). Using the incidence rates of EC, Golestan province was divided into high and low risk areas for EC (Figure 1). Data on incidence rates of EC were obtained from the Golestan population-based cancer registry. Mann-Whitney U test was used to assess the difference in levels of Se between the 2 areas. Pearson's correlation tests were used to assess the correlation between mean rice Se levels and the incidence rates of

EC in Golestan cities. A *p*-value of less than 0.05 was considered as significant. No human data or samples were collected in this study; therefore, there was no ethical consideration to be reviewed by our institutional ethical committee.

Results. In total, 69 rice samples were collected from different regions of Golestan province. Twenty-seven (39.1%) of the samples were collected from low EC-risk areas, and 42 (60.9%) from high EC-risk areas (Figure 1). An internal quality control of the analytic procedure was conducted throughout the study. Accuracy was evaluated by adding the standard Se (5 $\mu\text{g/l}$) into the sample, the mean recovery was 98.1% (93.7-105.4%). The coefficient of variation obtained was 2.8%. The maximum concentration of Se in rice samples was 0.5 mg/kg. The mean (SD) level of Se in rice samples in the low EC-risk area was 0.16 mg/kg (± 0.02), and in high EC-risk area was 0.35 mg/kg (± 0.02) ($p < 0.001$). We also found a significant correlation between rice Se levels and the incidence rate of EC in Golestan province (Pearson's correlation coefficient=0.65, $p=0.03$) (Figure 2).

Discussion. We found a relatively high concentration of Se in rice samples from the Golestan province of Iran (0.229 mg/kg). The mean Se content values in rice reported in literature were 0.020 mg/kg for Italy in 2007,¹⁰ and 0.05 mg/kg for Korea in 2004.¹¹ Our results also suggested that the level of Se in rice samples from high EC-risk areas was significantly




Figure 2 - Correlation between age standardized incidence rate (ASIR) of esophageal cancer and mean selenium (Se) concentrations in rice samples from Golestan cities, Iran.

higher than low risk areas. In some studies, Se has been investigated in relation to cancer risk due to its protective effects. Some studies have found Se status to be inversely associated with cancer risk. Negative associations have been found for various parameters of Se status and risks to cancers.¹²

Selenium was suggested as a possible factor in the regional variation of cancer mortality in China and other areas, specifically for EC.¹³ Similar results were reported on the levels of Se in soil samples from Golestan province.⁴ Semnani et al⁴ in 2010 reported a significantly higher level of soil Se in high EC-risk areas than low EC-risk areas and suggested soil Se level as a possible risk factor for EC in this region. Therefore, rice Se levels may be considered a possible contributor to the high incidence of EC in Golestan province.

The results of the present study were in line with those from previous studies,⁴ suggesting that contamination of the environment (soil, rice, and so forth) by xenobiotics including Se may be proposed as a possible risk factor for EC in this region. Therefore, it should be considered in designing cancer control programs in this area.

We found a significant correlation between the levels of Se in rice samples from Golestan province and the incidence rate of EC in this region. Soils contaminated by xenobiotics from aerial depositions and irrigation are likely to induce a corresponding contamination in harvested crops. Crops grown on contaminated land can carry pollution to humans or animals.¹⁴ However, farmers use chemical fertilizers for fertilization of products, and this may lead to the accumulation of xenobiotics in rice.¹⁵ Accordingly, one of the main routes for human exposure to xenobiotics is the soil-crop-food pathway.¹⁴ There may be a direct relationship between the composition of soil including xenobiotics and that of agricultural products such as rice. Therefore, environmental characteristics (for example, soil composition) should be closely monitored especially in high risk areas including Golestan province of Iran. Any change or abnormality should be considered an important health-related issue and should be addressed in health policy-making and control programs.

This study suffers from limitations of ecological designs including the ecologic fallacy of assumed equal exposure, and the absence of temporal relationship between exposure and outcome. So a causal relationship could not be concluded from our findings. Further individual-based studies are needed to determine the role of Se in the pathogenesis of EC in this region.

In conclusion, our results showed high levels of Se in rice samples and a significant positive relationship between rice Se levels and EC rates in the Golestan province of Iran. Considering the previously reported

correlation between soil Se levels and EC rate, it may be concluded that high soil and rice Se levels may play a possible role in the pathogenesis of EC in this area.

Received 31st October 2013. Accepted 1st April 2014.

From the Environmental Health Research Center (Rahimzadeh-Barzoki, Beirami), the Golestan Research Center of Gastroenterology and Hepatology (Joshiaghani, Semnani, Roshandel), Golestan University of Medical Sciences, Gorgan, and the Department of Environmental Health (Mansurian), Faculty of Health, Ilam University of Medical Sciences, Ilam, Iran. Address correspondence and re-prints request to: Dr. Gholamreza Roshandel, Number 77, Qaboosiehb Passage, Valiasr Street, PO Box 49166-53588, Gorgan, Iran. Fax: +98 (171) 2369210. E-mail: roshandel_md@yahoo.com

References

1. Ferlay J, Soerjomataram I, Ervik M, Dikshit R, Eser S, Mathers C, et al. GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC Cancer Base No. 11. Lyon (FR): International Agency for Research on Cancer; 2013. [Accessed 2013 July 11]. Available from: <http://globocan.iarc.fr>
2. Roshandel G, Sadjadi A, Aarabi M, Keshtkar A, Sedaghat SM, Nouraei SM, et al. Cancer incidence in Golestan province: report of an ongoing population-based cancer registry in Iran between 2004 and 2008. *Arch Iran Med* 2012; 15: 196-204.
3. Nouraei M, Pourshams A, Kamangar F, Sotoudeh M, Derakhshan MH, Akbari MR, et al. Ecologic study of serum selenium and upper gastrointestinal cancers in Iran. *World J Gastroenterol* 2004; 10: 2544-2546.
4. Semnani S, Roshandel G, Zendeabad A, Keshtkar A, Rahimzadeh H, Abdolahi N, et al. Soils selenium level and esophageal cancer: an ecological study in a high risk area for esophageal cancer. *J Trace Elem Med Biol* 2010; 24: 174-177.
5. Zhang B, Yang L, Wang W, Li Y, Li H. Environmental selenium in the Kaschin-Beck disease area, Tibetan Plateau, China. *Environ Geochem Health* 2011; 33: 495-501. Erratum in 2012; 34: 297-303.
6. Emmanuelle B, Virginie M, Fabienne S, Isabelle I, Martine PG, Bernard L, et al. Selenium exposure in subjects living in areas with high selenium concentrated drinking water: results of a French integrated exposure assessment survey. *Environ Int* 2012; 40: 155-161.
7. Appleton JD, Zhang Q, Green KA, Zhang G, Ge X, Liu X, et al. Selenium in soil, grain, human hair and drinking water in relation to esophageal cancer in the Cixian area, Hebei province, People's Republic of China *Applied Geochemistry* 2006; 21: 684-700.
8. Food Agriculture Organization. FAO statistical databases. Rome (IT): Food Agriculture Organization; 2004. [Access date 2013 August 25]. Available from: <http://www.fao.org/statistics/databases/en/>
9. Ashournia M, Aliakbar A. Determination of selenium in natural waters by adsorptive differential pulse cathodic stripping voltammetry. *J Hazard Mater* 2009; 168: 542-547.
10. Panigati M, Falciola L, Mussini P, Beretta G, Fancino RM. Determination of selenium in Italian rices by differential pulse cathodic stripping voltammetry. *Food Chemistry* 2007; 105: 1091-1098.
11. Choi Y, Kim J, Lee HS, Kim CI, Hwang IK, Park HK, et al. Selenium content in representative Korean foods. *Journal of Food Composition and Analysis* 2009; 22: 117-122.

12. Johnson CC, Fordyce FM, Rayman MP. Symposium on 'Geographical and geological influences on nutrition': Factors controlling the distribution of selenium in the environment and their impact on health and nutrition. *Proc Nutr Soc* 2010; 69: 119-132.
13. Chen J, Geissler C, Parpia B, Li J, Campbell TC. Antioxidant status and cancer mortality in China. *Int J Epidemiol* 1992; 21: 625-635.
14. Fu J, Zhou Q, Liu J, Liu W, Wang T, Zhang Q, et al. High levels of heavy metals in rice (*Oryza sativa L.*) from a typical E-waste recycling area in southeast China and its potential risk to human health. *Chemosphere* 2008; 71: 1269-1275.
15. Khaniki GJ, Zazoli MA. Cadmium and lead contents in rice (*Oryza sativa*) in the North of Iran. *International Journal of Agriculture & Biology* 2005; 7: 1026-1029.

Related Articles

Ge Z, Zhao C, Wang Y, Qian J. Cholecystectomy and the risk of esophageal and gastric cancer. *Saudi Med J* 2012; 33: 1073-1079.

Per H, Canpolat M, Sahin U, Gumus H, Konuskan B, Kumandas S. Serum and urine boron and selenium levels in children with resistant epilepsy. *Saudi Med J* 2012; 33: 942-947.

Ehteshamfar SM, Shapouri-Moghaddam A, Safarian M, Nematy M, Bahrami-Taghanaki H, Azizi H. Serum selenium concentration in Mashhad prisoners, Iran. *Saudi Med J* 2012; 33: 859-862.