

Molecular variation in *Leishmania* parasites from sandflies species of a zoonotic cutaneous leishmaniasis in northeast of Iran

Mitra Sharbatkhori^{1,2}, Adel Spotin^{3,4}, Heshmatollah Taherkhani², Mona Roshanghalb³ & Parviz Parvizi³

¹Laboratory Science Research Centre, Golestan University of Medical Sciences, Gorgan; ²Department of Parasitology and Mycology, School of Medicine, Golestan University of Medical Sciences, Gorgan; ³Molecular Systematics Laboratory, Parasitology Department, Pasteur Institute of Iran, Tehran; ⁴Parasitology Department, Medical Faculty, Tabriz University of Medical Sciences, Tabriz, Iran

ABSTRACT

Background & objectives: In the well-known zoonotic cutaneous leishmaniasis (ZCL) focus in Turkmen Sahara, border of Iran and Turkmenistan, ZCL has increased among humans in the past five years. The present study was undertaken to incriminate vectors of ZCL in the region, and to find molecular variation in *Leishmania* parasites.

Methods: The sandflies were sampled using CDC light-traps and sticky papers. All the sandflies were identified using morphological characters of the head and abdominal terminalia. DNA was extracted from the dissected thorax and attached anterior abdomen of individual female sandfly. *Leishmania* detection and identification of sandflies were performed using PCR, digestion of *Bsu*RI restriction enzyme and sequencing of ITS-rDNA gene and also by semi-nested PCR to amplify minicircle kinetoplast (k) DNA of *Leishmania*.

Results: *Leishmania* infections were detected in 26 out of 206 female sandflies. Of the infected sandflies, 18 were *Phlebotomus papatasi* while eight were *P. caucasicus/P. mongolensis*. Two infections of *L. turnica* were detected, one in *P. papatasi* and other in *P. caucasicus/P. mongolensis* and the rest of the sandflies were found infected with *L. major*.

Conclusion: Our finding showed that *L. major* had low diversity with only one common haplotype (GenBank Access No. EF413075). The novel haplotypes were discovered in *L. major* (GenBank Access No. KF152937) and in *L. turanica* (GenBank Access No. EF413079) in low frequency. These *Leishmania* parasites are circulating to maintain infections in the *P. papatasi* and *P. caucasicus/P. mongolensis* in Turkmen Sahara.

Key words Iran; ITS ribosomal DNA; *Leishmania major*; *L. turanica*; *Phlebotomus papatasi*; *P. caucasicus/P. mongolensis*

INTRODUCTION

Zoonotic cutaneous leishmaniasis (ZCL) is endemic disease in Iran and *Leishmania major* is the causative agent transmitted by sandflies¹⁻². Turkmen Sahara district, in northeast of Iran, border of Iran and Turkmenistan, is well-known and one of the most important endemic area of ZCL²⁻³. The haematophagous females of *Phlebotomus papatasi* (Scopoli) (Diptera: Psychodidae) have been incriminated as the main vector of the parasitic protozoan *L. major* (Yakimoff and Schokhor) (Kinetoplastida: Trypanosomatidae), the principal agent of ZCL⁴⁻⁵. In some ZCL endemic parts of Iran, as well as *P. papatasi*, other species of phlebotomine sandflies and reservoir hosts, like the gerbil *Rhombomys opimus* (Licht.) (Rodentia: Gerbillidae), have been found infected with *L. major* and unidentified *Leishmania*^{2,5-6}. Nevertheless, most *Leishmania* infections have been detected in *P. papatasi* followed by *P. (Paraphlebotomus) caucasicus* Marzinowsky and / or *P. mongolensis* Sinton which are likely to be the main and secondary vectors in Iran^{5,7}.

Phlebotomus caucasicus and *P. mongolensis* are frequently found in the burrows of the great gerbil, *R. opimus* and other reservoir hosts in Iran. The females of these sandfly species are similar and can not be separated morphologically and molecularly^{3, 8-9}.

Our study sites are part of an important ZCL focus in Golestan province, in northeast of Iran (Fig. 1). Recently, according to Disease Control Unit of the Health Centre report, ZCL cases increased significantly and became as an epidemiological problem over the past five years in some locations in Turkmen Sahara¹⁰. The first objective was to determine whether any molecular variation exists in *Leishmania* parasites. The second objective was to develop our knowledge regarding the detection of one or more *Paraphlebotomus* sandflies in which the *Leishmania* parasites survive and continue their transmission cycles. The last, was evaluation and finding new approaches of controlling surveillance of leishmaniasis in Turkmen Sahara focus. In addition, we tried to identify and type molecular *Leishmania* species in *P. papatasi* and *P. caucasicus* as main and potential vectors of ZCL



Map not to scale

Fig. 1: Locations of Gonbad-e Kâvus, Aqqala and Marveh Tapeh districts in Turkmen Sahara, Golestan province of Iran, where sandflies were sampled and screened for *Leishmania* infections.

in Turkmen Sahara in northeast of Iran. Some of our preliminary data has already been published in Persian in a regional journal¹¹.

The internal transcribed spacers (ITS) and kinetoplast (k) DNA of *Leishmania* parasites were selected to firmly identify and confirm the molecular analysis using nested and semi-nested PCR, respectively. Also restriction fragment length polymorphism (RFLP) and sequencing of ITS gene were performed for final verification^{12–15}.

MATERIAL & METHODS

Sandfly collections, identification and extraction of DNA

Sandflies were sampled from 15 villages of Turkmen Sahara, Golestan province, in northeast of Iran. Villages of our study sites are coordinated in Gonbad-e Qâbus or Gonbad-e Kâvus (37° 15' 00" N 55° 10' 02" E), Aqqala (37° 00' 50" N 54° 27' 18" E), and Marveh Tapeh (37° 54' 15" N 55° 57' 21" E) in a ZCL focus of this region (Fig. 1). Sandflies collected were from animal shelters, gerbil burrows and inside the houses in 2009 and 2010¹⁶ using CDC light-traps and sticky papers. All the sandflies were identified using morphological characters of the head and abdominal terminalia^{17–18}. DNA was extracted from the thorax and attached anterior abdomen of each sandfly as previously described¹⁹. Nested PCR of ITS-rDNA was used for detecting and identifying *Leishmania* species in sandflies.

A total of 206 female sandflies from these collections were screened for infections of *Leishmania* species by nested PCR. The PCR assay was carried out as pre the protocol described by Parvizi and Ready⁴. The first PCR was performed using forward primer IR1 reverse primer

IR2. The nested PCR was done using forward primer ITS1F and reverse primer ITS2R4⁴.

A semi-nested PCR of minicircle kinetoplast (k) DNA for identifying Leishmania species in sandflies

The primers LINR4 (forward), LIN17 (first-step reverse) and LIN19 (second-step reverse) were used for a semi-nested PCR¹⁹. The primers anneal within the conserved area of the minicircle and are also based on the Conserved Sequence Blocks recognized by Brewster *et al*²⁰.

Restriction fragment length polymorphism (RFLP) of ITS-rDNA gene

The sequence of standard strain (*L. major*, MHOM/SU/73/5ASKH; *L. infantum*, MHOM/TN/80/IPT1; *L. turanica*, MRHO/MN/83/MNR1 and *L. gerbilli*, MRHO/UZ/87/KD-87555) which have been registered in GenBank were used for sequencing analysis to select the suitable enzyme for digesting the PCR product in RFLP method.

Leishmania species were analysed using CLC DNA Workbench 5.2 software (CLC bio A/S, Aarhus, Denmark) to select the suitable enzyme which has different cut sites in different species of *Leishmania*, after sequencing standard strains of it. *BsuRI* (*HaeIII*) was selected with cut site GG↓CC as an appropriate enzyme for PCR product digestion. Endonuclease digestion was performed in a volume 30 µl include PCR product 10 µl, *BsuRI* (*HaeIII*) enzyme (Fermentas) 2 µl, 10× buffer 2 µl and distilled water 16 µl for 4 h at 37°C.

Furthermore, standard strains of *L. major*, *L. tropica* and *L. infantum* were used as positive controls. One of two negative controls was without restriction enzyme and the other one had no PCR product. After digesting the PCR product by endonuclease-restriction enzyme using *BsuRI* (*HaeIII*) enzyme, the fragments were analysed by using electrophoresis on 3% agarose gel containing ethidium bromide and ladder 50 bp (Fermentas).

The DNA fragments amplified by PCR were fractionated on 1.5% agarose gels, together with standard DNA fragments to permit sizing. PCR products were directly sequenced to identify *Leishmania* haplotypes infecting individual female sandflies, and all haplotypes were identified to species by phylogenetic analysis. For this, DNA sequences were edited and aligned using Sequencher™ 4.4.1 software, and the multiple alignments of new DNA haplotypes and homologous GenBank sequences were exported into MEGA software for phylogenetic analysis. The homologous GenBank sequences were selected and molecular phylogenetic tree was constructed for the Old World *Leishmania* species.

Table 1. *Leishmania* species identified in *P. papatasi* and *P. caucasicus/P. mongolensis* in different villages based on habitats and abdomens position using size of kDNA, RFLP of ITS-rDNA fragment and sequences

Location	Data collection										<i>P. caucasicus/P. mongolensis</i>										RFLP															
	Total (+)ve					Abdomen position					Habitat					Total (+)ve					Abdomen position					Habitat					<i>P. papatasi</i>					
	[kDNA]		(ITS)			FF	G	SG	IH	RB	ASH	FF	G	SG	IH	RB	ASH	FF	G	SG	IH	RB	ASH	FF	G	SG	IH	RB	ASH	L. major	L. turanica	L. major	L. turanica			
Gonbade-Kavous	Dash boron	70	(12)	[12]	13	10	47	(12)	0	7	63	(12)	6	(2)	[2]	0	3	(2)	3	1	3	(2)	2	11	1	3	(2)	2	1	2	0	0				
	Torshakli	24	(2)	[2]	0	0	24	(2)	0	0	24	(2)	4	(2)	[2]	0	1	3	(2)	0	1	(1)	3	(1)	2	0	0	1	1	1	1	1				
	Shurdgesh	5	(2)	[2]	0	0	5	(2)	0	0	5	(2)	1	0	0	1	0	1	0	0	1	0	0	2	0	0	0	0	0	0	0	0	0			
	Dozloom	11	(1)	[1]	0	0	11	(1)	0	0	11	(1)	2	0	0	1	1	0	0	0	1	1	0	1	0	0	0	0	0	0	0	0	0	0		
	Qeshlagh	1			0	0	1		0	0	1		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Aqqala	Daneshmand	20			0	1	19		0	1	19		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	Inche boron	3			0	2	1		1	2	0		2	0	0	2	0	2	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Okhitapeh	5			0	1	4		0	5	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Jafar bay	2			0	2	0		0	2	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Bibishirvan	1			0	1	0		0	1	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Marveh Tapeh	Farghesarepaen	8			0	0	8		0	0	1		2	0	0	2	0	0	2	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Gharagol	1	(1)	[1]	0	1	(1)	0	0	6	3	(1)	10	(4)	[4]	3	(1)	6	(3)	1	0	9	(4)	1	1	0	0	0	0	0	0	0	0	0	0	0
	Shorjepaen	3			0	0	3		0	0	8		1	0	0	3	1	0	3	1	0	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Souzes	2			0	0	2		1	1	0		6	2	2	2	2	2	2	2	2	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Kheyrkhajojliya	2			0	0	2		1	1	0		1	0	0	1	0	1	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	168	(18)	[18]	13	28	(1)	87	(17)	3	26	135	(18)	38	(8)	[8]	5	(1)	21	(5)	12	(2)	6	25	(7)	17	1	7	(1)	7	1	7	1	1	1		

FF— Full fed; G—Gravid; SG—Semi-gravid; IH—Inside houses; RB—Rhombomys burrows; ASH—Animal shelters.

RESULTS

Sandflies collection, Leishmania infections of sandflies identified by the nested PCR of ITS1-5.8 SrRNA gene-ITS2

Sandflies were widespread in September 2009 and 2010. *Phlebotomus papatasi* was abundant and predominant species in the region. The collections included the putative vectors of *L. major* causing ZCL in the region and in Iran (Table 1). The females of *P. caucasicus* and *P. mongolensis* could not be separated morphologically but, based on the presence of the males, between these two species, most were *P. caucasicus*^{8, 21-22}. Altogether, 206 female sandflies of two species were collected from 15 villages and screened for *Leishmania* infections. A total of 26 infections of *Leishmania* were discovered (18 in *P. papatasi* and 8 in *P. caucasicus/P. mongolensis*) (Table 1). For the detection of *Leishmania* infections, the female sandflies were captured and identified using ITS-5.8S rRNA-ITS2 gene. Two *Leishmania* infections were unidentified using RFLP. All the 26 independently ITS-rDNA fragments from each species were directly sequenced, to confirm their homology with species-specific GenBank sequences³. Haplotype TSH61 (GenBank Access No. EF413075) was identified in some GenBank sequences from strains of *L. major* originating from different locations including Iran, and it predominated in Iranian sandflies infected with this species. New haplotype TSH94 (GenBank Access No. KF152937) of *L. major* was also found in three infections.

Two infections of *L. turanica* were detected in one *P. papatasi* and other in *P. caucasicus/P. mongolensis* (Table 1). Both single *P. papatasi* and *P. caucasicus/P. mongolensis* sandflies with *L. turanica* sequences had the same haplotype TSH91 (GenBank Access No. EF413079). Haplotype TSH91 had previously been isolated from *R. opimus* from Mongolia and some Republics of the ex-USSR and also in Iran²³.

Leishmania infections of sandflies identified by the semi-nested PCR of minicircle kinetoplast (k) DNA

Most of each minicircle kDNA molecule was amplified by the semi-nested PCR, as reported by Aransay *et al*¹⁵, the size of the fragment was diagnostic for *L. major* (650 bp) compared with three other Old World species of *Leishmania* (720 bp). For amplifying minicircle kinetoplast (k) DNA, all the 26 *Leishmania* positive for ITS-rDNA gene were tried using the semi-nested PCR. Size specificity was 100% for 24 fragments of *L. major* and two fragments for *L. turanica* (Table 1).

DISCUSSION

Our findings showed *L. major* and *L. turanica* parasite species maintain infections in the *P. papatasi* and *P. caucasicus/P. mongolensis* in Turkmen Sahara, but it is important to discover the complementary roles of the sandflies species transmitting each *Leishmania* species among the reservoir hosts and to people²⁴⁻²⁶. Out of 26, 24 infections in wild females of *P. papatasi* and *P. caucasicus/P. mongolensis* were identified as *L. major*, and infection rates were significantly higher in *P. papatasi* from domestic animal shelters (18/18) than in *P. papatasi* caught in gerbil burrows (0/18) and also inside the houses (0/18). But infection rates were higher for gerbil burrows in *P. caucasicus/P. mongolensis* (7/8) than in domestic animal shelters (1/8) and no infection was detected in sandflies inside the houses (0/8) ($p < 0.01$). Perhaps *P. caucasicus/P. mongolensis* prefer to feed blood of rodents and rest in gerbil burrows²⁷⁻²⁹.

These two habitats are often <300 m apart in the Turkmen Sahara focus of ZCL, but its peridomestic populations have rarely been screened for *Leishmania* and few infections have been previously reported^{20, 26, 30}. Sandflies caught in gerbil burrows, contain with most infections needs in the burrows of the reservoir hosts, from where some infected females later disperse to domestic animal shelters⁸.

Nested PCR of ITS-rDNA have 100% specificity based on sequencing the amplified DNA fragments^{5, 31}. The ITS-rDNA gene was used for phylogenetic analysis of the *Leishmania* parasites from which the sequences originated based on a previous study in Iran and elsewhere³². The ITS-rDNA sequences obtained from the *P. mongolensis/P. caucasicus* and *P. papatasi* showed a similarity to the sequences deposited in GenBank. The results of a detailed phylogenetic analysis including characterization of such an approach could serve as a foundation for understanding the life cycle of *Leishmania* in sandflies (Fig. 2).

RFLP of (*Bsu*RI [*Hae*III] enzyme) ITS-rDNA gene was diagnostic for *L. major* to compare two mammalian *Leishmania* (*L. turanica* and *L. gerbilli*)¹³, because of the size of the amplified DNA fragment for *L. major* (650 bp) compared with three other species of *L. donovani*, *L. infantum* and *L. tropica* (720 bp)¹⁶. Semi-nested PCR of minicircle kDNA has the practical advantage of being diagnostic. There is no report for sizing fragment of minicircle kDNA for *L. turanica*. Based on sequencing the amplified DNA fragments the size of the fragment the visualized obtained bands for *L. turanica* which was equal to 720 bp similar to the standard strains of *L. donovani*, *L. infantum* and *L. tropica*²¹.




Fig. 2: Unrooted neighbour-joining tree showing the relationships of the haplotypes of the ITS1-5.8SrRNA gene fragment for the isolates of *Leishmania* species using PAUP.

None of *P. papatasi* and *P. causicus*/*P. mongolensis* in the present study had double infections unlike *R. opimus* in the ex-USSR and also in Iran. In Iran, several studies were carried out and found different *Leishmania* species in various parts of Iran but most of their work was based on conventional methods and some investigators employed molecular methods but they stopped after running and getting band in Agarose gel^{30, 32–34}.

Regarding our findings in this study, because of limited flight range, sandflies transmission of leishmaniasis within ZCL focus in Turkmen Sahara is often discontinuous, with characteristically small and separate foci close to the reservoir host habitats^{29, 35}. Also, infection rates were higher in semi-gravid and gravid than in females with red blood meals and for those without blood meals or eggs. Female sandflies need blood for egg production, some of the flies without blood or eggs might have been nulliparous and, therefore, had no chance to become infected^{36–37}.

The morphological and molecular similarity of female of *P. causicus* and *P. mongolensis* prevents a direct investigation of their roles in maintaining transmission of *L. major*^{9, 24} and other *Leishmania* species of gerbils^{5, 7}.

Now we can conclude that these sandfly species might be the vectors of *L. major* and *L. turanica* in human and/or in reservoir hosts of ZCL in Turkmen Sahara. In the future, more work should be carried out to test the status of sandflies as vectors of *L. major* and *L. turanica*.

ACKNOWLEDGEMENTS

The authors thank Mehdi Baghban for his help in the field work and Elnaz Alaee Novin for help in Molecular Systematics Laboratory. A part of this research was assigned by Mona Roshanghalb, M.Sc. student at Pasteur Institute of Iran, Tehran. The work was supported by the Pasteur Institute of Iran, grant number 367 awarded to Dr Parviz Parvizi. The collections of sandflies were made possible by the assistance of the Centre of Health Service in Gonband and Maraveh Tapeh, Golestan province, Iran. Authors thank Elham Esmaili, for language editing.

REFERENCES

1. Parvizi P, Alaee Novin E, Kazerooni PA, Ready PD. Low diversity of *Leishmania* parasites in sandflies and the absence of the great gerbil in foci of zoonotic cutaneous leishmaniasis in Fars province, southern Iran. *Trans R Soc Trop Med Hyg* 2013; 107: 356–62.
2. Rouhani S, Mirzaei A, Spotin A, Parvizi P. Novel identification of *Leishmania major* in *Hemichinus auritus* and molecular detection of this parasite in *Meriones libycus* from important foci of zoonotic cutaneous leishmaniasis in Iran. *J Infect Public Health* 2014; doi: 10.1016/j.jiph. 2013.12.002.
3. Mirzaei A, Rouhani S, Taherkhani H, Farahmand M, Kazemi B, Hedayati M, et al. Isolation and detection of *Leishmania* species among naturally infected *Rhombomys opimus*, a reservoir host of zoonotic cutaneous leishmaniasis in Turkmen Sahara, northeast of Iran. *Exp Parasitol* 2011; 129: 375–80.
4. Parvizi P, Ready PD. Nested PCRs and sequencing of nuclear ITS-rDNA fragments detect three *Leishmania* species of gerbils in sandflies from Iranian foci of zoonotic cutaneous leishmaniasis. *Trop Med Int Health* 2008; 13: 1159–71.
5. Killick-Kendrick R, Leaney A, Peters W, Rioux J, Bray R. Zoonotic cutaneous leishmaniasis in Saudi Arabia: The incrimination of *Phlebotomus papatasi* as the vector in the Al-Hassa oasis. *Trans R Soc Trop Med Hyg* 1985; 79: 252–5.
6. Mirzaei A, Rouhani S, Kazerooni PA, Farahmand M, Parvizi P. Molecular detection and conventional identification of *Leishmania* species in reservoir hosts of zoonotic cutaneous leishmaniasis in Fars Province, south of Iran. *Iranian J Parasitol* 2013; 8: 280–1.
7. Parvizi P, Benlarbi M, Ready PD. Mitochondrial and *Wolbachia* markers for the sandfly *Phlebotomus papatasi*: Little population

- differentiation between peridomestic sites and gerbil burrows in Isfahan province, Iran. *Med Vet Entomol* 2003; 17: 351–62.
8. Parvizi P, Taherkhani H, Ready PD. *Phlebotomus caucasicus* and *Phlebotomus mongolensis* (Diptera: Psychodidae): Indistinguishable by the mitochondrial cytochrome b gene in Iran. *Bull Entomol Res* 2010; 100: 415–20.
 9. Theodor O, Mesghali A. On the Phlebotominae of Iran. *J Med Entomol* 1964; 1: 285–300.
 10. Bordbar A, Parvizi P. High infection frequency, low diversity of *Leishmania major* and first detection of *Leishmania turanica* in human in northern Iran. *Acta Trop* 2014; 133: 69–74.
 11. Roshanghalb M, Parvizi P. Isolation and determination of *Leishmania major* and *Leishmania turanica* in *Phlebotomus papatasi* main vector of zoonotic cutaneous leishmaniasis in Turkemen Sahara Golestan province. *Mazand Univ Med Sci* 2012; 22: 74–83 (Persian).
 12. Bousslimi N, Aoun K, Ben-Abda I, Ben-Alaya-Bouafif N, Raouane M, Bouratbine A. Epidemiologic and clinical features of cutaneous leishmaniasis in southeastern Tunisia. *Am J Trop Med Hyg* 2010; 83: 1034–9.
 13. Cupolillo E, Grimaldi JG, Momen H, Beverley S. Intergenic region typing (IRT): A rapid molecular approach to the characterization and evolution of *Leishmania*. *Mol Biochem Parasitol* 1995; 73: 145–55.
 14. Mauricio I, Stothard J, Miles M. *Leishmania donovani* complex: Genotyping with the ribosomal internal transcribed spacer and the mini-exon. *Parasitol* 2004; 128: 263–7.
 15. Aransay AM, Scoulica E, Tselentis Y. Detection and identification of *Leishmania* DNA within naturally infected sandflies by seminested PCR on minicircle kinetoplastic DNA. *Appl Environ Microbiol* 2000; 66: 1933–8.
 16. Sudia W, Chamberlain R. Battery-operated light trap, an improved model. *Mosq News* 1962; 22: 126–9.
 17. Lewis D. A taxonomic review of the genus *Phlebotomus* (Diptera: Psychodidae). *Bull Br Mus* 1982; 45: 121–209.
 18. Nadim A, Javadian E. Key for species identification of sandflies (Diptera: Phlebotominae) of Iran. *J Pub Health* 1976; 5: 35–44.
 19. Parvizi P, Mauricio I, Aransay AM, Miles MA, Ready PD. First detection of *Leishmania major* in peridomestic *Phlebotomus papatasi* from Isfahan province, Iran: Comparison of nested PCR of nuclear ITS ribosomal DNA and semi-nested PCR of minicircle kinetoplast DNA. *Acta Trop* 2005; 93: 75–83.
 20. Brewster S, Aslett M, Barker DC. Kinetoplast DNA minicircle database. *Parasitol Today* 1998; 14: 437–8.
 21. Depaquit J, Ferté H, Léger N. The subgenus *Paraphlebotomus* (Phlebotomus—Phlebotominae—Psychodidae—Diptera): A review. Morphological and molecular studies (in French). *Ann Pharm Fr* 2000; 58: 333–40.
 22. Moin-Vaziri V, Depaquit J, Yaghoobi-Ershadi M. Variation between different geographical populations of *Phlebotomus (Paraphlebotomus) caucasicus* (Diptera: Psychodidae) in Iran. *Bull Soc Pathol Exot* 2007; 100: 211–5.
 23. Strelkova MV, Eliseev LN, Ponirovsky EN, Dergacheva TI, Annacharyeva DK, Erokhin PI, *et al.* Mixed leishmanial infections in *Rhombomys opimus*: A key to the persistence of *Leishmania major* from one transmission season to the next. *Ann Trop Med Parasitol* 2001; 95: 811–9.
 24. Killick-Kendrick R. Phlebotomine vectors of the leishmaniasis: A review. *Med Vet Entomol* 1990; 4: 1–24.
 25. Killick-Kendrick R, Ward R. Ecology of *Leishmania*. *Parasitology* 1981; 82: 143–52.
 26. Tabbabi A, Ghrab J, Aoun K, Ready PD, Bouratbine AD. Habitats of the sandfly vectors of *Leishmania tropica* and *L. major* in a mixed focus of cutaneous leishmaniasis in southeast Tunisia. *Acta Trop* 2011; 119: 131–7.
 27. Ali-Akbarpour M, Mohammadbeigi A, Tabatabaee S, Hatam G. Spatial analysis of eco-environmental risk factors of cutaneous leishmaniasis in southern Iran. *J Cutan Aesthet Surg* 2012; 5: 30–5.
 28. González C, Wang O, Strutz SE, González-Salazar C, Sánchez-Cordero V, Sarkar S. Climate change and risk of leishmaniasis in North America: Predictions from ecological niche models of vector and reservoir species. *PLoS Negl Trop Dis* 2010; 4: e585.
 29. Ready PD. Leishmaniasis emergence and climate change. *Rev Sci Tech* 2008; 27: 399–412.
 30. Oshaghi MA, Rasolian M, Shirzadi MR, Mohtarami F, Doosti S. First report on isolation of *Leishmania tropica* from sandflies of a classical urban cutaneous leishmaniasis focus in southern Iran. *Exp Parasitol* 2010; 126: 445–50.
 31. Schönian G, El Fari M, Lewin S, Schweynoch C, Presber W. Molecular epidemiology and population genetics in *Leishmania*. *Med Microbiol Immunol* 2001; 190: 61–3.
 32. Abai M, Rassi Y, Imamian H, Fateh M, Mohebbali M, Rafizadeh S, *et al.* PCR based on identification of vectors of zoonotic cutaneous leishmaniasis in Shahrood district, central of Iran. *Pak J Biol Sci* 2007; 10: 2061–5.
 33. Mohebbali M, Edrissian G, Shirzadi M, Akhouni B, Hajjaran H, Zarei Z, *et al.* An observational study on the current distribution of visceral leishmaniasis in different geographical zones of Iran and implication to health policy. *Travel Med Infect Dis* 2011; 9: 67–74.
 34. Sharifi I, Poursmaelian S, Aflatoonian MR, Ardakani RF, Mirzaei M, Fekri AR, *et al.* Emergence of a new focus of anthroponotic cutaneous leishmaniasis due to *Leishmania tropica* in rural communities of Bam district after the earthquake, Iran. *Trop Med Int Health* 2011; 16: 510–3.
 35. Ready PD. Biology of Phlebotomine sandflies as vectors of disease agents. *Ann Rev Entomol* 2013; 58: 227–50.
 36. Jafari R, Najafzadeh N, Sedaghat MM, Parvizi P. Molecular characterization of sandflies and *Leishmania* detection in main vector of zoonotic cutaneous leishmaniasis in Abarkouh district of Yazd province, Iran. *Asian Pac J Trop Med* 2013; 6: 792–7.
 37. Parvizi P, Alaeenovin E, Mohammadi S, Baghban N. Occurrence of low density of *Leishmania infantum* in sandflies from a new focus of visceral leishmaniasis in northwest of Iran. *J Vector Borne Dis* 2013; 50: 127–32.

Correspondence to: Dr Parviz Parvizi, Molecular Systematics Laboratory, Parasitology Department, Pasteur Institute of Iran, Tehran, Iran.
E mail: parp@pasteur.ac.ir; parpparvizi@yahoo.com

Received: 21 August 2013

Accepted in revised form: 17 December 2013