

A Review on the Metabolic Disorders of Iodine Deficiency

Azad Reza Mansourian

Biochemistry and Metabolic Disorder Research Center, Gorgan Medical School,
Golestan University of Medical Sciences, Gorgan, Iran

Abstract: Iodine is in the crucial parts of two hormones of T4 and T3 produced by the thyroid glands which are essential for all the aspects of human metabolisms. It is demonstrated that iodine deficiency can be considered as sole cause of many thyroid abnormalities including mental disorders. Iodine deficiency of sufficient degree to cause hypothyroidism during fetus life and early infancy will be accompanied with brain abnormality possibly to the stage of mental retardation. The iodine deficiency among subjects in their early stage of childhood is not as severe as those in their fetus or infancy. In adult subjects the severe iodine deficiency can be also associated with mental disorders due to the direct side effects of hypothyroidism occurred by lack of iodine. The clinical manifestation of iodine deficiency show itself with psychological disorders in adult subjects. The status of iodine within blood can be evaluated through measurement of urinary iodine level and the low urinary concentration is an indicative of hypothyroidism. Mental retardation and brain damage due to iodine deficiency can be prevented if iodine supplementation prescribed duly on time.

Key words: Hypothyroidism, infant, adult, mental retardation, brain damage

INTRODUCTION

Thyroid gland produces two important hormones tetra iodothyronin or thyroxine (T4) and triiodothyronine (T3). These two hormones are essential for human physiological process and metabolisms (Reinehr, 2010; Reinehr *et al.*, 2008; Mansourian *et al.*, 2008; Mansourian, 2010b-e).

Thyroid hormone deficiency accompanied with serious adverse effects and it will have a catastrophic consequences particularly among infants and young children. In some circumstances, the physical and mental status of infant both are affected. The biosynthesis of thyroid hormones exclusively depends on the presence of iodine which is entirely thyroid based element and the thyroid is the sole site of iodine absorption and storage (Mansourian, 2011a; Mansourian *et al.*, 2007; Mansourian, 2010e).

The oxidized iodine within thyroid is transferred on the tyrosyle residues of thyroglobulin and converted into tyrosyle iodinated substances monoiodo and diiodo tyrosyle (MIT, DIT) on thyroglobuline. The coupling process followed between MIT and DIT, DIT and DIT produce T3 and T4, respectively still on the thyroglobuline molecules. It is therefore, interestingly obvious that 4 and 3 in T4 and T3 are designated for the number of iodine atoms present in the biochemical structure of thyroid hormones and this is why the iodine reside in crucial center of thyroid hormones and in fact T4,

T3 without iodine atom are only amino acids derivatives. Thyroglobuline is hydrolyzed subsequently by the lysosome enzymes within the extra follicular space of thyroid and eventually T4 and T3 are secreted into blood circulation (Mansourian, 2011a).

Iodine therefore, reside in the central core of manufacturing T4 and T3 in thyroid and iodine deficiency in human eventually lead to serious metabolic disorders. Iodine deficiency caused tremendous headaches for health care providers world wide which can be occurred from early stage of fetus life (Zimmermann and Andersson, 2011; Delange, 2002; Laurberg *et al.*, 2006; Kapil, 2010; Andersson *et al.*, 2010; Rajatanavin, 2007; Kurtoglu *et al.*, 2004; Jaruratanasirikul *et al.*, 2006; Kung, 2007; Luton *et al.*, 2011; Delange *et al.*, 2002; Vern *et al.*, 2003). Iodine deficiency eventually lead to hypothyroidism, only in rare cases it can cause hyperthyroidism but it is not absolutely common (Pearce *et al.*, 2004; Haldimann *et al.*, 2005; El-Ghawi and Al-Sadeq, 2006; Thomson *et al.*, 2008; Chavasit *et al.*, 2002; Mansourian, 2010b, c, e).

The level of thyroid hormones dictate the amount of iodine for the production of T4, T3. The iodine level also depends on the age, physiological status of each individual (Glinoe, 2007, 2003, 1999, 2001; Luton *et al.*, 2011; Delange, 1994; Laurberg *et al.*, 2000, 2001, 2007; Zarei *et al.*, 2009).

It seems that pregnancy, fetus life, infancy and childhood, are the periods with highest requirement of

iodine, respectively due to higher level of thyroid hormones which are duly required (Sidibeel, 2007; Glinoyer, 2007, 1999; Delange, 1998). Various studies indicated the vital and crucial requirement of iodine intake for the proper amount of thyroid hormone to prevent particularly the brain damage during fetus and infancy. It is therefore, absolutely necessary to have enough iodine supplementation during pregnancy and in early stage of infancy and childhood to prevent the occurrence of mentally retarded newborns (Delange, 1998; Glinoyer, 2007, 2003, 1999; Eftekhari *et al.*, 2007).

The role of iodine in fetus life and early infancy: Iodine is absorbed into thyroid gland through an active transport mediated by Na/K pump which itself regulated and stimulated by Thyroid Stimulating Hormone (TSH) a pituitary hormone. The situation of iodine deficiency seems to be even worse during pregnancy, due to mental and physical growth adverse effects. It is widely advised that pregnant women dietary regiment should be supplemented with enough iodine to prevent possible hypothyroidism which can happen due the iodine shortage (Berbel *et al.*, 2007; Zimmermann, 2007a; Maberly, 1994). It should be mentioned that in addition to iodine deficiency, any other metabolic disorders which occur genetically on the biochemical structure of thyroid hormone translocator protein and receptor which are present on the cell surface and nucleus of T4 and T3 target tissues also can have a similar adverse effects, because in either cases thyroid hormones messages are not relayed into the tissues (Glinoyer, 1999, 2001, 2007). In addition to mental and physical disturbances iodine deficiency of large extend to cause hypothyroidism will interfere with the physiological status of pregnancy accompanied with catastrophic event in some cases and fetus death eventually (Berbel *et al.*, 2007; Zimmermann, 2007a; Maberly, 1994). It should be also mentioned that the stage at which iodine deficiency occur can play an important role in the presentation of signs and symptoms of iodine deficiency during pregnancy. The first and the early stage of second trimester of pregnancy and early infancy is an utmost importance in regard to iodine deficiency and much damage to the fetus occur during this phase of pregnancy due to onset of maternal hypothyroidism in first and early second trimester of pregnancy (DeLong *et al.*, 1985; Vulmsa *et al.*, 1989; Christensen and Davis, 2004). Under any condition iodine deficiency accelerate the rate of hypothyroidism in maternal life up to 10-12 times, therefore, to prevent the side effects of fetus damage due to maternal iodine deficiency maternal hypothyroidism should be checked out regularly to prevent any possible mental and physical

retardation of fetus, although the newborns should be assessed for their blood thyroid hormone shortly after delivery to prevent similar consequences (Delange, 1997; Glinoyer, 1997).

The role of iodine deficiency in brain tissues: In case of iodine deficiency the anatomy of brain tissues are not properly formed, the cell production and differentiation altered due to iodine deficiency. Iodine deficiency enough to cause hypothyroidism during fetus life and early infancy eventually cause mental disturbance with possible mental retardation (Eayrs and Taylor, 1951; Madeira *et al.*, 1988a, b). Thyroid hormone also play an important roles in nervous system metabolic pathway, therefore it seems the lack of iodine if converted to hypothyroidism can eventually leave the newborn with mental abnormality (Ambrogini *et al.*, 2005; Desouza *et al.*, 2005; Lemkine *et al.*, 2005; Montero-Pedrazuela *et al.*, 2006; Emden and Jack, 2011; Jaruratanasirikul *et al.*, 2006; Rajatanavin, 2007).

The metabolic pathway of thyroid hormones within the brain follow almost the same pathways to other tissues and it is the free thyroid hormones within blood circulation which play an important role in regulating metabolic pathways. It means T4 and T3 following biosynthesis within thyroid gland secreted into blood circulation, transported by transporting protein which are synthesized in the liver, reach the target tissues and bind to the receptors following passage through cell membrane by thyroid hormone translocator protein. Like other peripheral tissues in brain, T4 can also be converted into T3 which is a more potent hormone with different speed to accommodate itself with brain actual requirement (Escobar-Morreale *et al.*, 1995, 1999; Mansourian, 2011a, b; Mansourian 2010a, c).

T4 and T3 play an utmost important role in growing fetus physiological life and particularly in the brain formation during the first trimester of pregnancy which fetus completely depends to the maternal thyroid hormone also the requirement is decreased as fetus grow further but it is a slow process which need extra attention to have normal mental and physical status (Kurtoglu *et al.*, 2004; De Escobar *et al.*, 1987, 1990, 1985, 2004; Obregon *et al.*, 1984; Porterfield and Hendrich, 1992; Woods *et al.*, 1984; Contempre *et al.*, 1993; Bernal and Pekonen, 1984; Vulmsa *et al.*, 1989).

Malnutrition may lead to iodine deficiency hypothyroidism: One of main problem of thyroid disorder end-up with hypothyroidism is iodine deficiency due to the nutritional status and the deficiency of iodine in the dietary regiment. A high portion of thyroid deficiency

leading to the clinical manifestation of cretinism among newborn and children are due to maternal dietary iodine deficiency during pregnancy and early phase of infancy which is also caused by the fact that in many region of the world the nutrient used as daily dietary consumption are iodine deficient as well and the iodine supplementation in such region is a urgent matter to be considered (Zimmermann and Andersson, 2011; Delange, 2002; Zhong *et al.*, 1983; Hetzel and Hay, 1979; McMichael *et al.*, 1980). Experimentally hypothyroidism are produced during pregnancy by omitting the iodine from daily dietary. These studies demonstrated the whole clinical manifestation of thyroid hormone deficiency on the growing fetus following iodine deletion from nutritional regiment. The other studies indicated that the scenario was reversed, by iodine supplementation (Kapil, 2010; Laurberg *et al.*, 2006, 2001; Potter *et al.*, 1981).

Although the fetus physical and mental growth in early phase of pregnancy entirely depends to the maternal circulation of thyroid hormones but experimental evidence, proved that in latter stage of pregnancy also the growing fetus itself is able to produce thyroid hormones but the hormones should be supplemented also by maternal circulation to have a fetus with normal mental and physical growth (McMichael *et al.*, 1980; Hetzel *et al.*, 1988; Delange, 1998; Glinoe, 1999; Mansourian, 2010e, d; Shahmohammadi *et al.*, 2008; Mansourian *et al.*, 2010a, b; Mansourian *et al.*, 2007; Marjani *et al.*, 2008; Mansourian and Ahmadi, 2010).

It should be kept in mind that although thyroid hormone or iodine supplementation may eventually improve physical growth but the fetus and newborn mental disorder, caused by T4 and T3 deficiency, might be an irreversible pathway (Auso *et al.*, 2004; Koibuchi and Chin, 2000; Chan and Kilby, 2000; Delange, 2000, 2001).

The iodine deficiency associated with various abnormalities ranging from pregnancy disorder affecting the fetus with various anomalies and in some cases, accompanied with fatal outcome (Glinoe, 2007; Sidibeel, 2007; Delange, 1994, 1998; Andersson *et al.*, 2010; Zimmermann, 2007a, 2009; Mansourian, 2010e). Even following giving birth to the child by iodine deficient mother the newborn can be accompanied with possible risk of mental retardation. The iodine deficiency during adulthood up to the elderly is accompanied with goiter and associated consequences of hypothyroidism. Accordingly iodine related malnutrition, malabsorption and related nutritional status in any stage of life can be accompanied with hypothyroidism adverse effects (Auso *et al.*, 2004; Koibuchi and Chin, 2000; Chan and Kilby, 2000; Delange, 2000, 2001).

The mechanism behind brain damage during fetus and early life due to iodine deficiency relay on the fact that, lack of iodine eventually accompanied with hypothyroidism which will be resulted in the reduction of T4 and T3 serum concentration, followed by adverse alteration of gene expression particularly within the brain and nerves tissues, of fetus and neonate (Auso *et al.*, 2004; Koibuchi and Chin, 2000; Chan and Kilby, 2000; Delange, 2000, 2001).

The reversibility of brain damage is not certain, by thyroid hormone, it is this dilemma which manifest the importance of iodine supplementation during pregnancy and early stage of infancy which is absolutely crucial particularly for the mental function (Auso *et al.*, 2004; Koibuchi and Chin, 2000; Chan and Kilby, 2000; Delange, 2000, 2001).

As it was mentioned above the different stage of pregnancy is of utmost importance for the fetus mental growth (Dobbing and Sands, 1973). Following termination of pregnancy and in early phase of infancy, the brain still is not developed properly and during both stages of fetus life and early infancy the presence of thyroid hormones, crucially are required and lack of iodine prevent or at least slow the process of brain formation into proper shape. It is vital to remember that the brain development in the early stage of fetal life entirely relay on the maternal circulation of enough thyroid hormone but as fetus grows further it is less dependent on the maternal thyroid hormones circulation (Dobbing and Sands, 1973). Under any circumstance, the lack of iodine which lead to the thyroid hormone deficiency may eventually ended with mentally retard infant (Dobbing and Sands, 1973). It should be mentioned that in some cases iodine excess is accompanied with adverse effects and following iodine overload the signs and symptoms of hypothyroidism and hyperthyroidism are both can be exhibited and this is why iodine intake should be only on physician and nutritionists recommendation (Carceller *et al.*, 2011; Kurtoglu *et al.*, 2009; Khashu *et al.*, 2005; Ergur *et al.*, 2011; Teng *et al.*, 2011).

The significant role which played by the fetal and maternal T4 and T3, prove that the clinical care should be taken during pregnancy on the amount of iodine supplementation to prevent any thyroid abnormalities. Also fetus can produce thyroid hormone to substitute the maternal thyroid hormones but one should remember, that the circulating maternal thyroid hormones always support the fetus thyroid hormones requirements particularly during first half of pregnancy and it seems maternal thyroid hormone partly required up to the very last days of pregnancy (Zimmermann, 2007b; Zimmermann, 2009;

Bernal and Pekonen, 1984; Kooistra *et al.*, 2006; Vulsma *et al.*, 1989). The gene expression which occurred through the combination of T4 and T3 to their nuclear receptor is considered as a cornerstone to our understanding in how thyroid hormone facilitates their effects in producing various proteins required for brain development. Although thyroid hormones are binding to their receptor, within nuclear region of brain cell but the free hormone reaches the receptor through a passage, assisted by a translocation protein present on the brain cell membrane (Delange *et al.*, 1989).

Various degrees of iodine deficiency and Brain development during pregnancy: Iodine deficiency can be subdivided into severe, mild and moderate, if severe iodine deficiency was demonstrated it can be accompanied by serious physical and mental disturbance during fetus life. There are controversial arguments on condition of mild and moderate iodine deficiency but pregnant women possible hypothyroidism due to iodine deficiency may eventually cause the fetus physical and mental disturbances but this area of studies is not absolutely clear and it is matter for further investigations (Kooistra *et al.*, 2006; Vulsma *et al.*, 1989; Delange *et al.*, 1989).

Under any condition iodine deficiency during fetal life may lead to fatal outcome and studies showed the iodine supplementation reduce the severity of scenario and such circumstance are reversed on iodine prescription. Various studies from different part of the world indicated that the fatal outcome of iodine deficiency in infancy also reduced following iodine supplementation (Pharoah and Connolly, 1987; Cobra *et al.*, 1997; Thilly *et al.*, 1980, 1986; DeLong *et al.*, 1997; Dobbing, 1974).

The iodine supplementation in such circumstance, enable the thyroid to produce enough T4 and T3 for the proper functioning and stimulation of healthy brain development. The throughout studies confirmed that iodine deficiency is not only belong to the poor and developing countries, but the iodine deficiency and possible brain damage can be observed even in well developed countries but it should be emphasized that the most adverse effects of iodine deficiency are seen among infant and it is why higher iodine requirement is necessary for younger population due to their higher metabolic rate and growth developments (Zimmermann and Andersson, 2011; Delange, 2002, 1990; Dobbing, 1974).

Iodine deficiency among children: The iodine deficiency among subjects in their early stage of childhood, is not severe as those in their fetus or neonatal life. It has been

shown that cognition of children are not adversely affected but it has to be mentioned that the iodine supplementation enhance the intelligence ability of children (Aghini-Lombardi *et al.*, 1995; Santiago-Fernandez *et al.*, 2004). Other studies indicated the clinical manifestation have been improved to some acceptable range after iodine supplementation among children with iodine deficiency (Zimmermann *et al.*, 2006). The intelligence capacity of those children, that their fetus life were associated with severe iodine deficient mother was lower, compared to those children that their fetus life was accompanied with moderate maternal iodine deficiency (Fierro-Benitez *et al.*, 1974; Kochupillai *et al.*, 1986; Huda *et al.*, 1999). Under any circumstances, some reports indicated the iodine deficiency of even moderate to severe cases can be accompanied by reducing the intellectual capacity among children to some extent (Bleichrodt and Born, 1994; Qian *et al.*, 2005). In addition to the earlier reported studies, other investigation indicated that mild to moderate iodine deficiency cause the reducing psychomotor and mental development, neuromuscular disorders, reducing learning capability, slowing psychomotor development and many similar results which mostly related with the level iodine deficiency compared with the control subjects. It has also been demonstrated that iodine supplementation in some instance enhance the rate of intelligence and psychomotor ability (Bleichrodt *et al.*, 1989; Vermiglio *et al.*, 1990; Fenzi *et al.*, 1990; Tiwari *et al.*, 1996; Shresta, 1994; Van den Briel *et al.*, 2000; Aghini-Lombardi *et al.*, 1995).

In spite of all documented reports on the positive role played by iodine supplementation, on the cognitive and psychomotor abilities but this area of research are controversially discussed (Zimmermann *et al.*, 2007; Zimmermann, 2009; Mansourian, 2010c; Mansourian *et al.*, 2010b; Mansourian and Ahmadi, 2010).

Iodine deficiency among adults population: It has been reported that iodine deficiency of enough degree to be considered as a severe, can profoundly associated with mental disorder due to the direct side effects of thyroid hypo function and subsequent overt hypothyroidism with possible, cerebral abnormalities in adult population. The iodine deficiency, eventually, show itself with psychological disorder, to such extent that daily life and routine procedure of adults subjects can be effected negatively (Hollowell and Hammon, 1997; Mansourian *et al.*, 2007; Mansourian, 2010c; Mansourian, 2011a). Although iodine deficiency do not demonstrate a fatal outcome among adults but the adult iodine deficiency is associated with problems, that can adversely interfere with routine procedure of daily life

and normal metabolism (Aghini-Lombardi *et al.*, 1999; Vanderpump *et al.*, 1995; Laurberg *et al.*, 1998; Szabolcs *et al.*, 1997; Garcia-Mayor *et al.*, 1999; Mansourian *et al.*, 2008; China, Marjani *et al.*, 2008; Mansourian, 2010b).

In rare cases iodine supplementation trigger hyperthyroidism: Surprisingly, iodine supplementation programs in some cases, stimulate the thyroid gland towards hyperthyroidism. The concept behind this observation, derived from this fact that the iodine deficiency itself contribute the thyroid cells proliferation, thyrocyte and Thyroid Stimulating Hormone (TSH) receptor mutation. TSH is a pituitary hormone which stimulate thyroid gland to produce T4 and T3 after binding to the TSH Receptor (TSHR) on the thyroid gland. The thyroid gland begin to produce T4 and T3 following TSH binds to TSHR (Ergur *et al.*, 2011; Carceller *et al.*, 2011; Kurtoglu *et al.*, 2009; Delange *et al.*, 1999; Baltisberger *et al.*, 1995; Mansourian, 2011a).

It seems, that in some from of moderate to sever iodine deficiency iodine supplementation possibly stimulated the iodine-induced hyperthyroidism but this is a pathway which can be reversed eventually (Dremier *et al.*, 1996).

The other possibilities of excessive iodine supplementation, is the adverse effect of such supplementation, on inducing the immune system to be become activated against thyrocyte in genetically susceptible individual. In some other studies, the iodine supplementation stimulate the auto-immunity against the thyroid and possible production of auto antibodies against thyroid gland (Jonckheer *et al.*, 1992; Delange and Lecomte, 2000; McConahey *et al.*, 1962; Mansourian, 2010a).

The iodine supplementation of excessive amount even in healthy subjects, can stimulate the healthy thyroid into a condition in which the immune system activate against the peroxidase enzyme responsible for iodine peroxidation prior to T4 and T3 production within the thyroid gland. This observation which is adversely affect the thyroid, is reversed into normal condition eventually. The findings of this area of research are not and globally homogenously are agreed (Kahaly *et al.*, 1998; Hollowell *et al.*, 2002; Braverman, 1994; Pearce *et al.*, 2002; Yang *et al.*, 2007; Teng *et al.*, 2006; Mansourian, 2010e).

There are also some reports indicating that if excessive amount of iodine taken for a long period of times the iodine overload hypothyroidism, can be a eventual outcome, of such undesired dietary regiments. It is also reported that iodine excess to the level which

stimulate the thyroid into goiter condition is done either by iodine stimulation of auto immunity against thyroid gland or the excess iodine and the eventual absorption by the thyroid, lead to the temporary inhibition of T4 and T3 synthesis through the Wolff- Chikoff effect, in those individual which are prone to such adverse effect. (Roti *et al.*, 1981; Suzuki *et al.*, 1965; Tonglet *et al.*, 1992; Knudsen *et al.*, 2000a).

As whole the adverse effects of iodine excess on the thyroid gland are of both hypo and hyperthyroid disorder as explained above. It seems that the iodine deficiency and iodine overload both can interfere with proper mechanism of thyroid gland and in susceptible subjects either of iodine deficiency and excess can possibly accompanied with side effects of thyroid disorders (McGuire and Galloway, 1994).

How circulating blood T4 and T3 do their physiological function in the targets tissues: Most of T4 and T3 are transported through blood circulation by proteins which are synthesized within the liver (Mansourian, 2011b) and the free hormone which are the true active hormones pass through target tissues via a translocator protein complex resided on the membrane of target tissues. In addition to what it was mentioned the T4 and T3 nuclear receptor is shown to play a crucially important passage in this mechanism for the function of thyroid hormone in the brain in fetus and early infancy (Laterra and Goldstein, 2000; Dumitrescu *et al.*, 2004; Friesema *et al.*, 2004, 2005; Perez-Castillo *et al.*, 1985; Schwartz and Oppenheimer, 1978; Strait *et al.*, 1990).

Any physiological modification to this system can also cause some abnormalities to the tissues involved and if the brain and nervous system associated with this above mentioned disorder the fetus and infant manifest a sever clinical symptoms resulting from hypothyroidism due to iodine deficiency.

The thyroid hormones do their physiological function by regulation which take place through genetical manipulation to give the satisfactory speed to the various biochemical reactions of the brain and other tissues, through related enzymes of a specified reaction at specific genes. It should be mentioned that in general the exact scenario in how T4 and T3 behave to exercise their influence on gene manipulation particularly during fetus life is a subject for further investigation in this area of medical science and major portion of our knowledge in this region is through experimental research on animal models (Schwartz *et al.*, 1997; Krebs *et al.*, 1996; Sampson *et al.*, 2000; Dowling *et al.*, 2001; Sinha *et al.*, 2008; Farsetti *et al.*, 1992; Zou *et al.*, 1994). The present experimental data on how T4 and T3 influence gene

manipulation indicate that during infancy the brain tissues are more prone towards thyroid hormones and it seems that the brain development in this stage of life do happen through the gene expression regulated by thyroid hormones. The thyroid hormone clearly play an important role in regulation of adults psychological status up to the late stage of life and it is demonstrated that hormone therapy to treat hypothyroidism reverses the patient condition to the large extend (Diez *et al.*, 2008; Bauer *et al.*, 2002; Simon *et al.*, 2002; Baumgartner *et al.*, 1994; Bauer *et al.*, 2005).

Serum iodine evaluation can be assessed through urinary iodine measurement: Iodine within blood circulation can be assessed through, measurement of urinary iodine-concentration, the low level of urinary concentration is a indicative of hypothyroidism (Mansourian *et al.*, 2007; Mansourian, 2010e; Shahmohammdi *et al.*, 2008).

The simple goiter is a clinical manifestation explaining the possibility of reduced amount of intake iodine (Mansourian, 2010b, c). The TSH and Thyroglobulin (Tg), are also two indicators which can demonstrate the physiological status of thyroid gland with potential indication of iodine status.

TSH, measurement and its negative correlation with thyroid hormones of T4 and T3 clearly can give a clue to the thyroid function tests. In iodine deficiency the Tg which is an thyroid follicular space protein is released within blood circulation and its measurement can be an indicative iodine status of human body. The combination of TSH and Tg measurement are useful indicative for the assessment of iodine concentration (Spencer and Wang, 1995; Knudsen *et al.*, 2001; Mansourian, 2011a).

The estimation of thyroid hormone of T4 and T3 can not give a prices clue to the iodine level and should not be used precisely to assess the iodine status. TSH serum level physiologically is controlled by the amount of T4 and T3 but on the other hand the level of T4 and T3 may in some way reflect the amount of iodine and alternatively the TSH measurement can in some cases also be an indicative of iodine level. It should be mentioned that TSH can not be thoroughly trusted at different age groups to estimate serum iodine level but its concentration most probably can be used as serum iodine status. Due to the utmost importance of thyroid hormones and iodine assessment in early infancy TSH assessment as single test seems to be a valuable probe into in how newborn thyroid function at crucial days of life. It is therefore, suggested that serum T4 and T3 level in infants can ultimately be examined by TSH measurements at a time when the newborns require the optimal thyroid hormones

level to prevent brain damage (Andersen *et al.*, 2008; Knudsen *et al.*, 2000b; Jaruratanasirikul *et al.*, 2009; Jaruratanasirikul *et al.*, 2006).

CONCLUSION

- Thyroid gland produce two important hormones, of tetra idothyronin or thyroxine (T4) and triiodothyronine (T3). These hormones are essential for all the aspect of human physiological process and metabolisms
- Iodine deficiency to the degree leading to hypothyroidism during fetus life and early infancy causing the brain malfunction to the stage which brain tissues are not properly formed with possible mental retardation outcome
- The iodine deficiency among subjects in their early stage of childhood is not as severe as those in their fetus or neonatal life. It has been shown that cognition of children are not adversely affected but it has to be mentioned that the iodine supplementation enhance the intelligence ability of children
- It has been reported that iodine deficiency of enough degree to be considered as a sever, can profoundly associated with mental disorder in adult population due to the direct side effects of overt hypothyroidism with possible, cerebral abnormalities. The clinical manifestation of iodine deficiency eventually show itself with psychological disorders, to the such extend that the daily life and routine procedure of adults subjects can be effected seriously
- The status of iodine within blood can be evaluated through measurement of urinary iodine level and the low urinary concentration is a indicative of hypothyroidism. Some of the side effects of thyroid malfunction in such conditions can be corrected particularly among adults subjects by iodine supplementation if it is done on time.

REFERENCES

- Aghini-Lombardi, F., A. Pinchera, L. Antonangeli, T. Rago and L. Chiovato *et al.*, 1995. Mild iodine deficiency during fetal/ neonatal life and neuropsychological impairment in Tuscany. *J. Endocrinol. Invest.*, 18: 57-62.
- Aghini-Lombardi, F., L. Antonangeli, E. Martino, P. Vitti and D. Maccherini *et al.*, 1999. The spectrum of thyroid disorders in an iodine-deficient community: The Pescoporgano survey. *J. Clin. Endocrinol. Metab.*, 84: 561-566.

- Ambrogini, P., R. Cuppini, P. Ferri, C. Mancini and S. Ciaroni *et al.*, 2005. Thyroid hormones affect neurogenesis in the dentate gyrus of adult rat. *Neuroendocrinology*, 81: 244-253.
- Andersen, S., J. Karmisholt, K.M. Pedersen and P. Laurberg, 2008. Reliability of studies of iodine intake and recommendations for number of samples in groups and in individuals. *Br. J. Nutr.*, 99: 813-818.
- Andersson, M., I. Aeberli, N. Wust, A.M. Piacenza and T. Bucher *et al.*, 2010. The swiss iodized salt program provides adequate iodine for school children and pregnant women but weaning infants not receiving iodine-containing complementary foods as well as their mothers are iodine deficient. *J. Clin. Endocrinol. Metab.*, 95: 5217-5224.
- Auso, E., R. Lavado-Autric, E. Cuevas, F.E. Del-Rey, G.M. De Escobar and P. Berbel, 2004. A moderate and transient deficiency of maternal thyroid function at the beginning of fetal neurogenesis alters neuronal migration. *Endocrinology*, 145: 4037-4047.
- Baltisberger, B.L., C.E. Minder and H. Burgi, 1995. Decrease of incidence of toxic nodular goiter in a region of Switzerland after full correction of mild iodine deficiency. *Eur. J. Endocrinol.*, 132: 546-549.
- Bauer, M., A. Heinz and P.C. Whybrow, 2002. Thyroid hormones, serotonin and mood: of synergy and significance in the adult brain. *Mol. Psychiatry*, 7: 140-156.
- Bauer, M., E.D. London, N. Rasgon, S.M. Berman and M.A. Frye *et al.*, 2005. Supraphysiological doses of levothyroxine alter regional cerebral metabolism and improve mood in bipolar depression. *Mol. Psychiatry*, 10: 456-469.
- Baumgartner, A., M. Bauer and R. Hellweg, 1994. Treatment of intractable non-rapid cycling bipolar affective disorder with high-dose thyroxine: An open clinical trial. *Neuropsychopharmacology*, 10: 183-189.
- Berbel, P., M.J. Obregón, J. Bernal, F. Escobar del Rey and G. Morreale de Escobar, 2007. Iodine supplementation during pregnancy: A public health challenge. *Trends Endocrinol. Metab.*, 18: 338-343.
- Bernal, J. and F. Pekonen, 1984. Ontogenesis of the nuclear 3,5,3-triiodothyronine receptor in the human fetal brain. *Endocrinology*, 114: 677-679.
- Bleichrodt, N., F.E.D. Rey, G.M.D. Escobar, I. Garcia and C. Rubio, 1989. Iodine Deficiency. Implications for Mental and Psychomotor Development in Children. In: *Iodine and the Brain*, DeLong, G.R., J. Robbins and P.G. Condliffe (Eds.). Plenum Press Publication, New York, pp: 269-287.
- Bleichrodt, N. and M.P. Born, 1994. A Metaanalysis of Research on Iodine and its Relationship to Cognitive Development. In: *The Damaged Brain of Iodine Deficiency*, Stanbury, J.B. (Ed.). Cognizant Communication Publication, New York, pp: 195-200.
- Braverman, L.E., 1994. Iodine and the thyroid: 33 years of study. *Thyroid*, 4: 351-356.
- Carceller, D.B., P.L. Segundo, P.N. Beltran, A.S. Paris, P.C. Hernandez, M.M. Villanueva and D.A. Olivares, 2011. Short-term results of treatment with (131)I in patients with multinodular goiter: Effect of the associated degree of hyperthyroidism and other variables. *Rev. Esp. Med. Nucl.*
- Chan, S. and M.D. Kilby, 2000. Thyroid hormone and central nervous system development. *J. Endocrinol.*, 165: 1-8.
- Chavasit, V., P. Malaivongse and K. Judprasong, 2002. Study on stability of iodine in iodated salt by use of different cooking model conditions. *J. Food Comp. Anal.*, 15: 265-276.
- Christensen, V.L. and G.S. Davis, 2004. Maternal dietary iodide influences turkey embryo thyroid function. *Int. J. Poult. Sci.*, 3: 550-557.
- Cobra, C., Muhilal, K. Rusmil, D. Rustama and Djatnika *et al.*, 1997. Infant survival is improved by oral iodine supplementation. *J. Nutr.*, 127: 574-578.
- Contempre, B., E. Jauniaux, R. Calvo, D. Jurkovic, S. Campbell and G.M. de Escobar, 1993. Detection of thyroid hormones in human embryonic cavities during the first trimester of pregnancy. *J. Clin. Endocrinol. Metabol.*, 77: 1719-1722.
- De Escobar, G.M., R. Pastor, M.J. Obregon and F.E. del Rey, 1985. Effects of maternal hypothyroidism on the weight and thyroid hormone content of rat embryonic tissues, before and after onset of fetal thyroid function. *Endocrinology*, 117: 1890-1900.
- De Escobar, G.M., M.J. Obregon and F.E. del Rey, 1987. Fetal and maternal thyroid hormones. *Hormone Res.*, 26: 12-27.
- De Escobar, G.M., R. Calvo, M.J. Obregon and F.E. del Rey, 1990. Contribution of maternal thyroxine to fetal thyroxine pools in normal rats near term. *Endocrinology*, 126: 2765-2767.
- De Escobar, G.M., M.J. Obregon and F.E. Del-Rey, 2004. Role of thyroid hormone during early brain development. *Eur. J. Endocrinol.*, 151: U25-U37.
- DeLong, G.R., J.B. Stanbury and R. Fierro-Benitez, 1985. Neurological signs in congenital iodine-deficiency disorder (endemic cretinism). *Dev. Med. Child Neurol.*, 27: 317-324.

- Delange, F., J. De Vijlder, G.M. de Escobar, P. Rochiccioli and S. Varrone, 1989. Significance of Early Diagnostic Data in Congenital Hypothyroidism: Report of the Subcommittee on Neonatal Hypothyroidism of the European Thyroid Association. In: Research in Congenital Hypothyroidism, Delange, F., D.A. Fisher and D. Glinoe (Eds.). Plenum Press, New York, pp: 225-234.
- Delange, F., 1990. Iodine Nutrition and Risk of Thyroid Irradiation from Nuclear Accidents. In: Iodine Prophylaxis following Nuclear Accidents, Rubery, E. and E. Smales (Eds.). Pergamon Press, Oxford, pp: 45-53.
- Delange, F., 1994. The disorders induced by iodine deficiency. *Thyroid*, 4: 107-128.
- Delange, F., 1997. Neonatal screening for congenital hypothyroidism: Results and perspectives. *Hormone Res.*, 48: 51-61.
- Delange, F., 1998. Screening for congenital hypothyroidism used as an indicator of the degree of iodine deficiency and of its control. *Thyroid*, 8: 1185-1192.
- Delange, F., B. de Benoist and D. Alnwick, 1999. Risks of iodine-induced hyperthyroidism after correction of iodine deficiency by iodized salt. *Thyroid*, 9: 545-556.
- Delange, F., 2000. Endemic Cretinism. In: The Thyroid. A Fundamental and Clinical Text, Braverman, L.E. and R.D. Utiger (Eds.). Lippincott, Philadelphia, PA., pp: 743-754.
- Delange, F. and P. Lecomte, 2000. Iodine supplementation: Benefits outweigh risks. *Drug Saf.*, 22: 89-95.
- Delange, F., 2001. Iodine deficiency as a cause of brain damage. *Postgrad. Med. J.*, 77: 217-220.
- Delange, F., 2002. Iodine deficiency in Europe and its consequences: An update. *J. Nucl. Med. Mol. Imaging.*, 29: S404-S416.
- Delange, F., H. Burgi, Z.P. Chen and J.T. Dunn, 2002. World status of monitoring iodine deficiency disorders control programs. *Thyroid*, 12: 915-924.
- Delong, G.R., P.W. Leslie, S.H. Wang, X.M. Jiang and M.L. Zhang *et al.*, 1997. Effect on infant mortality of iodination of irrigation water in a severely iodine-deficient area of China. *Lancet*, 350: 771-773.
- Desouza, L.A., U. Ladiwala, S.M. Daniel, S. Agashe, R.A. Vaidya and V.A. Vaidya, 2005. Thyroid hormone regulates hippocampal neurogenesis in the adult rat brain. *Mol. Cell Neurosci.*, 29: 414-426.
- Diez, D., C. Grijota-Martinez, P. Agretti, G. De Marco and M. Tonacchera *et al.*, 2008. Thyroid hormone action in the adult brain: Gene expression profiling of the effects of single and multiple doses of triiodo-L-thyronine in the rat striatum. *Endocrinology*, 149: 3989-4000.
- Dobbing, J. and J. Sands, 1973. Quantitative growth and development of human brain. *Arch. Dis. Child.*, 48: 757-767.
- Dobbing, J., 1974. The later growth of the brain and its vulnerability. *Pediatrics*, 53: 2-6.
- Dowling, A.L., E.A. Iannacone and R.T. Zoeller, 2001. Maternal hypothyroidism selectively affects the expression of neuroendocrine-specific protein a messenger ribonucleic acid in the proliferative zone of the fetal rat brain cortex. *Endocrinology*, 142: 390-399.
- Dremier, S., F. Coppee, F. Delange, G. Vassart, J.E. Dumont and J. Van Sande, 1996. Thyroid autonomy: Mechanism and clinical effects. *J. Clin. Endocrinol. Metab.*, 81: 4187-4193.
- Dumitrescu, A.M., X.H. Liao, T.B. Best, K. Brockmann and S. Refetoff, 2004. A novel syndrome combining thyroid and neurological abnormalities is associated with mutations in a monocarboxylate transporter gene. *Am. J. Hum. Genet.*, 74: 168-175.
- Eayrs, J.T. and S.H. Taylor, 1951. The effect of thyroid deficiency induced by methylthiouracil on the maturation of the central nervous system. *J. Anat.*, 85: 350-358.
- Eftekhari, M.H., Z. Mazloom, A. Ahmadi and H.M. Khosravi, 2007. Body mass index and thyroid function in adolescent girls. *Pak. J. Biol. Sci.*, 10: 905-909.
- El-Ghawi, U.M. and A.A. Al-Sadeq, 2006. Determination of iodine in Libyan food samples using epithermal instrumental neutron activation analysis. *Biol. Trace Elem. Res.*, 111: 31-40.
- Ender, P.J. and M.M. Jack, 2011. Iodine-induced neonatal hypothyroidism secondary to maternal seaweed consumption: A common practice in some Asian cultures to promote breast milk supply. *J. Paediatr. Child Health*, 10.1111/j.1440-1754.2010.01972.x
- Ergur, A.T., O. Evliyaoglu, Z. Siklar, P. Bilir, G. Ocal and M. Berberoglu, 2011. Evaluation of thyroid functions with respect to iodine status and TRH test in chronic autoimmune thyroiditis. *J. Clin. Res. Pediatr. Endocrinol.*, 3: 18-21.
- Escobar-Morreale, H.F., M.J. Obregon, F.E. del Rey and G.M. de Escobar, 1995. Replacement therapy for hypothyroidism with thyroxine alone does not ensure euthyroidism in all tissues, as studied in thyroidectomized rats. *J. Clin. Invest.*, 96: 2828-2838.
- Escobar-Morreale, H.F., M.J. Obregon, F.E. del Rey and G.M. de Escobar, 1999. Tissue-specific patterns of changes in 3,5,3'-triiodo-L-thyronine concentrations in thyroidectomized rats infused with increasing doses of the hormone. Which are the regulatory mechanisms?. *Biochimie*, 81: 453-462.

- Farsetti, A., B. Desvergne, P. Hallenbeck, J. Robbins and V.M. Nikodem, 1992. Characterization of myelin basic protein thyroid hormone response element and its function in the context of native and heterologous promoter. *J. Biol. Chem.*, 267: 15784-15788.
- Fenzi, G.F., L.F. Giusti, F. Aghini-Lombardi, L. Bartalena and C. Marcocci *et al.*, 1990. Neuropsychological assessment in schoolchildren from an area of moderate iodine deficiency. *J. Endocrinol. Invest.*, 13: 427-431.
- Fierro-Benitez, R., I. Ramirez, E. Estrella and J.B. Stanbury, 1974. The Role of Iodine Deficiency in Intellectual Development in an Area of Endemic Goiter. In: *Endemic Goiter and Cretinism: Continuing Threats to World Health*, Dunn, J. and G.A. Medeiros-Neto, (Eds.). Pan American Health Organization (PAHO) Scientific Publ, Washington, pp: 135-142.
- Friesema, E.C., A. Grueters, H. Biebermann, H. Krude and A. von Moers *et al.*, 2004. Association between mutations in a thyroid hormone transporter and severe X-linked psychomotor retardation. *Lancet*, 364: 1435-1437.
- Friesema, E.C., J. Jansen, C. Milici and T.J. Visser, 2005. Thyroid hormone transporters. *Vitam. Horm.*, 70: 137-167.
- Garcia-Mayor, R.V., M. Rios, E. Fluiters, L.F. Mendez, E.G. Garcia-Mayor and A. Andrade, 1999. Effect of iodine supplementation on a pediatric population with mild iodine deficiency. *Thyroid*, 9: 1089-1093.
- Glinoer, D., 1997. The regulation of thyroid function in pregnancy: Pathways of endocrine adaptation from physiology to pathology. *Endocrine Rev.*, 18: 404-433.
- Glinoer, D., 1999. What happens to the normal thyroid during pregnancy. *Thyroid*, 9: 631-635.
- Glinoer, D., 2001. Pregnancy and iodine. *Thyroid*, 11: 471-481.
- Glinoer, D., 2003. Feto-maternal repercussions of iodine deficiency during pregnancy. An update. *Ann. Endocrinol. (Paris)*, 64: 37-44.
- Glinoer, D., 2007. The importance of iodine nutrition during pregnancy. *Public Health Nutr.*, 10: 1542-1546.
- Haldimann, M., A. Alt, A. Blanc and K. Blondeau, 2005. Iodine content of food groups. *J. Food Comp. Anal.*, 18: 461-471.
- Hetzel, B.S. and I.D. Hay, 1979. Thyroid function, iodine nutrition and fetal brain development. *Clin. Endocrinol.*, 11: 445-460.
- Hetzel, B.S., J. Chavadej and B.J. Potter, 1988. The brain in iodine deficiency. *Neuropathol. Applied Neurobiol.*, 14: 93-104.
- Hollowell, J.G. and W.H. Hannon, 1997. Teratogen update: Iodine deficiency, a community teratogen. *Teratology*, 55: 389-405.
- Hollowell, J.G., N.W. Staehling, W.D. Flanders, W.H. Hannon, E.W. Gunter, C.A. Spencer and L.E. Braverman, 2002. Serum TSH, T4 and thyroid antibodies in the United States population (1988 to 1994): National health and nutrition examination survey (NHANES III). *J. Clin. Endocrinol. Metab.*, 87: 489-499.
- Huda, S.N., M.S.G. McGregor, K.M. Rahman and A. Tomkins, 1999. Biochemical hypothyroidism secondary to Iodine deficiency is associated with poor school achievement and cognition in Bangladeshi children. *J. Nutr.*, 129: 980-987.
- Jaruratanasirikul, S., J. Chukamnerd, O. Koranantakul, P. Chanvitan, P. Ruaengrairatanaroj and H. Sriplung, 2006. The relationship of maternal iodine status and neonatal thyrotropin concentration: A study in Southern Thailand. *J. Pediatr. Endocrinol. Metab.*, 19: 727-732.
- Jaruratanasirikul, S., P. Sangsupawanich, O. Koranantakul, P. Chanvitan and P. Ruaengrairatanaroj *et al.*, 2009. Maternal iodine status and neonatal thyroid-stimulating hormone concentration: A community survey in Songkhla, Southern Thailand. *Public Health Nutr.*, 12: 2279-2284.
- Jonckheer, M.H., B. Velkeniers, L. Vanhaelst and M. Van Blerk, 1992. Further characterization of iodide-induced hyperthyroidism based on the direct measurement of intrathyroidal iodine stores. *Nucl. Med. Commun.*, 13: 114-118.
- Kahaly, G.J., H.P. Dienes, J. Beyer and G. Hommel, 1998. Iodide induces thyroid autoimmunity in patients with endemic goitre: A randomised, double-blind, placebo-controlled trial. *Eur. J. Endocrinol.*, 139: 290-297.
- Kapil, U., 2010. Successful efforts toward elimination iodine deficiency disorders in India. *Indian J. Community Med.*, 35: 455-468.
- Khashu, M., P. Chessex and J.P. Chanoine, 2005. Iodine overload and severe hypothyroidism in a premature neonate. *J. Pediatr. Surg.*, 40: E1-E4.
- Knudsen, N., E. Christiansen, M. Brandt-Christensen, B. Nygaard and H. Perrild, 2000a. Age and sex-adjusted iodine/creatinine ratio. A new standard in epidemiological surveys? Evaluation of three different estimates of iodine excretion based on casual urine samples and comparison to 24 h values. *Eur. J. Clin. Nutr.*, 54: 361-363.

- Knudsen, N., I. Bulow, T. Jorgensen, P. Laurberg, L. Ovesen and H. Perrild, 2000b. Comparative study of thyroid function and types of thyroid dysfunction in two areas in Denmark with slightly different iodine status. *Eur. J. Endocrinol.*, 143: 485-491.
- Knudsen, N., I. Bulow, T. Jorgensen, H. Perrild, L. Ovesen and P. Laurberg, 2001. Serum Tg-A sensitive marker of thyroid abnormalities and iodine deficiency in epidemiological studies. *J. Clin. Endocrinol. Metab.*, 86: 3599-3603.
- Kochupillai, N., C.S. Pandav, M.M. Godbole, M. Mehta and M.M.S. Ahuja, 1986. Iodine deficiency and neonatal hypothyroidism. *Bull. WHO*, 64: 547-551.
- Koibuchi, N. and W.W. Chin, 2000. Thyroid hormone action and brain development. *Trends Endocrinol. Metab.*, 4: 123-128.
- Kooistra, L., S. Crwaford, A.L. van Baar, E.P. Bruowres and V.J. Pop, 2006. Neonatal effects of maternal hypothyroxinemia during early pregnancy. *Pediatrics*, 117: 161-167.
- Krebs, J., R.L. Means and P. Honegger, 1996. Induction of calmodulin kinase IV by the thyroid hormone during the development of rat brain. *J. Biol. Chem.*, 271: 11055-11058.
- Kung, A.W., 2007. Iodine nutrition of pregnant and lactating women in Hong Kong, where intake is of borderline sufficiency. *Public Health Nutr.*, 10: 1600-1601.
- Kurtoglu, S., M. Akcakus, C. Kocaoglu, T. Gunes and N. Budak *et al.*, 2004. Iodine status remains critical in mother and infant in Central Anatolia (Kayseri) of Turkey. *Eur. J. Nutr.*, 43: 297-303.
- Kurtoglu, S., L. Akin, M.A. Akin and D. Coban, 2009. Iodine overload and severe hypothyroidism in two neonates. *J. Clin. Res. Pediatr. Endocrinol.*, 1: 275-277.
- Lattera, J. and G.W. Goldstein, 2000. Ventricular Organization of Cerebrospinal Fluid: Blood-brain Barrier, Brain Edema and Hydrocephalus. In: *Principles of Neural Science*, Kandel, E.R., J.H. Schwartz and T.M. Jessell (Eds.). McGraw-Hill, New York, pp: 1288-1301.
- Laurberg, P., K.M. Pedersen, A. Hreidarsson, N. Sigfusson, E. Iversen and P.R. Knudsen, 1998. Iodine intake and the pattern of thyroid disorders: a comparative epidemiological study of thyroid abnormalities in the elderly in Iceland and in Jutland, Denmark. *J. Clin. Endocrinol. Metab.*, 83: 765-769.
- Laurberg, P., S.B. Nohr, K.M. Pedersen, A.B. Hreidarsson and S. Andersen *et al.*, 2000. Thyroid disorders in mild iodine deficiency. *Thyroid*, 10: 951-963.
- Laurberg, P., I.B. Pedersen, N. Knudsen, L. Ovesen and S. Andersen, 2001. Environmental iodine intake affects the type of nonmalignant thyroid disease. *Thyroid*, 11: 457-469.
- Laurberg, P., T. Jorgensen, H. Perrild, L. Ovesen and N. Knudsen *et al.*, 2006. The Danish investigation on iodine intake and thyroid disease, DanThyr: Status and perspectives. *Eur. J. Endocrinol.*, 155: 219-228.
- Laurberg, P., S. Anderson, R.I. Bjarnadottir, A. Carle and A.B. Hreidarsson *et al.*, 2007. Evaluating iodine deficiency in pregnant women and young infants-complex physiology with a risk of misinterpretation. *Public Health Nutr.*, 10: 1547-1552.
- Lemkine, G.F., A. Raj, G. Alfama, N. Turque and Z. Hassani *et al.*, 2005. Adult neural stem cell cycling *in vivo* requires thyroid hormone and its α receptor. *FASEB J.*, 19: 863-865.
- Luton, D., C. Alberti, E. Vuillard, G. Ducarme, J.F. Oury and J. Guibourdenche, 2011. Iodine deficiency in northern Paris area: Impact on fetal thyroid mensuration. *PLoS One*, 6: e14707-e14707.
- Maberly, G.F., 1994. Iodine deficiency disorders: Contemporary scientific issues. *J. Nutr.*, 124: 1473S-1478S.
- Madeira, M.D., M. Paula-Barbosa, A. Cadete-Leite and M.A. Tavares, 1988a. Unbiased estimate of cerebellar granule cell numbers in hypothyroid and in sex-age-matched control rats. *J. Hirnforsch.*, 29: 587-594.
- Madeira, M.D., M. Paula-Barbosa, A. Cadete-Leite and M.A. Tavares, 1988b. Unbiased estimate of hippocampal granule cell numbers in hypothyroid and in sex-age-matched control rats. *J. Hirnforsch.*, 29: 643-650.
- Mansourian, A.R., E.O. Ghaemi, A.R. Ahmadi, A. Saifi, A.V. Moradi and S. Bakhshandehnosrat, 2007. A survey of urinary iodine concentration in South East of Northern Iran. *Pak. J. Biol. Sci.*, 10: 2166-2171.
- Mansourian, A.R., E. Ghaemi, A.R. Ahmadi, A. Marjani, A. Saifi and S. Bakhshandehnosrat, 2008. Serum lipid level alterations in subclinical hypothyroid patients in Gorgan (South East of Caspian Sea). *Chinese Clin. Med.*, 3: 206-210.
- Mansourian, A.R., 2010a. A review on hyperthyroidism: Thyrotoxicosis under surveillance. *Pak. J. Biol. Sci.*, 13: 1066-1076.
- Mansourian, A.R., 2010b. Metabolic pathways of tetraiodothyronine (T₄) and triiodothyronine (T₃) production by thyroid gland: A review of articles. *Pak. J. Biol. Sci.*, 14: 1-12.
- Mansourian, A.R., 2010c. The immune system which adversely alter thyroid functions: A review on the concept of autoimmunity. *Pak. J. Biol. Sci.*, 13: 765-774.

- Mansourian, A.R., 2010d. The state of serum lipids profiles in sub-clinical hypothyroidism: A review of the literature. *Pak. J. Biol. Sci.*, 13: 556-562.
- Mansourian, A.R., 2010e. Thyroid function tests during first-trimester of pregnancy: A review of literature. *Pak. J. Biol. Sci.*, 13: 664-673.
- Mansourian, A.R. and A.R. Ahmadi, 2010. Correlation between inverse age and serum thyroxine level among children and adolescents. *J. Clin. Diagn. Res.*, 4: 3196-3200.
- Mansourian, A.R., A.R. Ahmadi, A. Saifi and S. Bakhshandehnosrat, 2010a. The children reference range of thyroid hormones in Northern Iran. *Pak. J. Biol. Sci.*, 13: 862-865.
- Mansourian, A.R., A.R. Ahmadi, H.R. Mansourian, A. Saifi, A. Marjani, G.R. Veghari and E. Ghaemi, 2010b. Maternal thyroid stimulating hormone level during the first trimester of pregnancy at the South-East of the caspian sea in Iran. *J. Clin. Diagn. Res.*, 4: 2472-2477.
- Mansourian, A.R., 2011a. Abnormal serum thyroid hormones concentration with healthy functional gland: A review on the metabolic role of thyroid hormones transporter proteins. *Pak. J. Biol. Sci.*, 14: 313-326.
- Mansourian, A.R., 2011b. Central dogma in thyroid dysfunction: A review on structure modification of TSHR as a cornerstone for thyroid abnormalities. *Pak. J. Biol. Sci.*, 14: 170-181.
- Marjani, A., A.R. Mansourian, E.O. Ghaemi, A. Ahmadi and V. Khori, 2008. Lipid peroxidation in serum of hypothyroid patients (in Gorgan-South East of Caspian Sea). *Asian J. Cell Biol.*, 3: 47-50.
- McConahey, W.M., F.R. Keating Jr., O.H. Beahr and L.B. Woolner, 1962. On the increasing occurrence of Hashimoto's thyroiditis. *J. Clin. Endocrinol. Metab.*, 22: 542-544.
- McGuire, J. and R. Galloway, 1994. *Enriching Lives. Overcoming Vitamin and Mineral Malnutrition in Developing Countries.* World Bank Publication, Washington, DC.
- McMichael, A.J., J.D. Potter and B.S. Hetzel, 1980. Iodine Deficiency, Thyroid Function and Reproductive Failure. In: *Endemic Goiter and Endemic Cretinism: Iodine Nutrition in Health and Disease*, Stanbury, J.B. and B.S. Hetzel (Eds.). John Wiley and Sons Publication, New York, pp: 445-460.
- Montero-Pedrazuela, A., C. Venero, R. Lavado-Autric, I. Fernandez-Lamo, K.M. Garcia-Verdugo, J. Bernal and A. Guadano-Ferraz, 2006. Modulation of adult hippocampal neurogenesis by thyroid hormones: Implications in depressive-like behavior. *Mol. Psychiatry*, 11: 361-371.
- Obregon, M.J., J. Mallol, R. Pastor, G. Morreale de Escobar and F. Escobar del Rey, 1984. L-thyroxine and 3,3',5-triiodo-L-thyronine in rat embryos before onset of fetal thyroid function. *Endocrinology*, 114: 305-307.
- Pearce, E.N., A.R. Gerber, D.B. Gootnick, L.K. Khan, R. Li, S. Pino and L.E. Braverman, 2002. Effects of chronic iodine excess in a cohort of long-term American workers in West Africa. *J. Clin. Endocrinol. Metab.*, 87: 5499-5502.
- Pearce, E.N., S. Pino, X. He, H.R. Bazrafshan, S.L. Lee and L.E. Braverman, 2004. Sources of dietary iodine: Bread, cows' milk and infant formula in the Boston area. *J. Clin. Endocrinol. Metab.*, 89: 3421-3424.
- Perez-Castillo, A., J. Bernal, B. Ferreiro and T. Pans, 1985. The early ontogenesis of thyroid hormone receptor in the rat fetus. *Endocrinology*, 117: 2457-2461.
- Pharoah, P.O.D. and K.J. Connolly, 1987. A controlled trial of iodinated oil for the prevention of endemic cretinism: A long-term follow-up. *Int. J. Epidemiol.*, 16: 68-73.
- Porterfield, S.P. and C.E. Hendrich, 1992. Tissue iodothyronine levels in fetuses of control and hypothyroid rats at 13 and 16 days gestation. *Endocrinology*, 131: 195-200.
- Potter, B.J., G.H. McIntosh and B.S. Hetzel, 1981. The Effect of Iodine Deficiency on Fetal Brain Development in the Sheep. In: *Fetal Brain Disorders. Recent Approaches to the Problem of Mental Deficiency*, Hetzel, B.S. and R.M. Smith (Eds.). Elsevier/North Holland Biomedical Press Publication, Amsterdam, pp: 119-148.
- Qian, M., D. Wang, W.E. Watkins, V. Gebiski, Y.Q. Yan, M. Li and Z.P. Chen, 2005. The effects of iodine on intelligence in children: A meta-analysis of studies conducted in China. *Asia Pacific J. Clin. Nutr.*, 14: 32-42.
- Rajatanavin, R., 2007. Iodine deficiency in pregnant women and neonates in Thailand. *Public Health Nutr.*, 10: 1602-1605.
- Reinehr, A., A. Asa, G. De Sousa, R. Dieffenbach and W. Andler, 2008. Thyroid hormones and their relation to weight status. *Horm Res.*, 70: 51-57.
- Reinehr, T., 2010. Obesity and thyroid function. *Mol. Cell Endocrinol.*, 316: 165-171.
- Roti, E., A. Gnudi, L.E. Braverman, G. Robuschi and R. Emanuele *et al.*, 1981. Human cord blood concentrations of thyrotropin, thyroglobulin and iodothyronines after maternal administration of thyrotropin-releasing hormone. *J. Clin. Endocrinol. Metab.*, 53: 813-817.

- Sampson, D., M.R. Pickard, A.K. Sinha, I.M. Evans, A.J. Leonard and R.P. Ekins, 2000. Maternal thyroid status regulates the expression of neuronal and astrocytic cytoskeletal proteins in the fetal brain. *J. Endocrinol.*, 167: 439-445.
- Santiago-Fernandez, P., R. Torres-Barahona, J.A. Muela-Martinez, G. Rojo-Martinez and E. Garcia-Fuentes *et al.*, 2004. Intelligence quotient and iodine intake: A cross-sectional study in children. *J. Clin. Endocrinol. Metab.*, 89: 3851-3857.
- Schwartz, H.L. and J.H. Oppenheimer, 1978. Ontogenesis of 3,5,3'-triiodothyronine receptors in neonatal rat brain, dissociation between receptor concentration and stimulation of oxygen consumption by 3,5,3'-triiodothyronine. *Endocrinology*, 103: 943-948.
- Schwartz, H.L., M.E. Ross and J.H. Oppenheimer, 1997. Lack of effect of thyroid hormone on late fetal rat brain development. *Endocrinology*, 138: 3119-3124.
- Shahmohammdi, F., A.R. Mansourian and H.R. Mansourian, 2008. Serum thyroid hormone level in women with nausea and vomiting in early pregnancy. *J. Med. Sci.*, 8: 507-510.
- Shresta, R.M. 1994. Effect of iodine and iron supplementation on physical, psychomotor and mental development in primary schoolchildren in Malawi. Ph.D. Thesis, Wageningen Agricultural University, Wageningen, Netherlands.
- Sidibeel, H., 2007. Reflections on mental retardation and congenital hypothyroidism: Effects of trace mineral deficiencies. *Sante*, 17: 41-50.
- Simon, N.M., D. Blacker, N.B. Korbly, S.G. Sharma, J.J. Worthington, M.W. Otto and M.H. Pollack, 2002. Hypothyroidism and hyperthyroidism in anxiety disorders revisited: New data and literature review. *J. Affect. Disord.*, 69: 209-217.
- Sinha, R.A., A. Pathak, V. Mohan, S. Bandyopadhyay, L. Rastogi and M.M. Godbole, 2008. Maternal thyroid hormone: A strong repressor of neuronal nitric oxide synthase in rat embryonic neocortex. *Endocrinology*, 149: 4396-4401.
- Spencer, C.A. and C.C. Wang, 1995. Thyroglobulin measurement. Techniques, clinical benefits and pitfalls. *Endocrinol. Metab. Clin. North Am.*, 24: 841-863.
- Strait, K.A., H.L. Schwartz, A. Perez-Castillo and J.H. Oppenheimer, 1990. Relationship of c-erbA mRNA content to tissue triiodothyronine nuclear binding capacity and function in developing and adult rats. *J. Biol. Chem.*, 265: 10514-10521.
- Suzuki, H., T. Higuchi, K. Sawa, S. Ohtaki and Y. Horiuchi, 1965. Endemic coast goitre in Hokkaido, Japan. *Acta Endocr. (Copenh)*, 50: 161-176.
- Szabolcs, I., J. Podoba, J. Feldkamp, O. Dohan and I. Farkas *et al.*, 1997. Comparative screening for thyroid disorders in old age in areas of iodine deficiency, long-term iodine prophylaxis and abundant iodine intake. *Clin. Endocrinol.*, 47: 87-92.
- Teng, W., Z. Shan, X. Teng, H. Guan and Y. Li *et al.*, 2006. Effect of iodine intake on thyroid diseases in China. *N. Engl. J. Med.*, 354: 2783-2793.
- Teng, X., Z. Shan, Y. Chen, Y. Lai and J. Yu *et al.*, 2011. hypothyroidism and autoimmune thyroiditis: A cross-sectional study based on two chinese communities with different iodine intake levels. *Eur. J. Endocrinol.*,
- Thilly, C.H., R. Lagasse, G. Roger, P. Bourdoux and A.M. Ermans, 1980. Impaired Fetal and Postnatal Development and High Perinatal Death-Rate in a Severe Iodine Deficient Area. In: *Thyroid Research VIII*, Stockigt, J.R., S. Nagataki, E. Meldrum, J.W. Barlow and P.E. Harding (Eds.). Australian Academy of Science, Canberra, Australia, pp: 20-23.
- Thilly, C.H., P.P. Bourdoux and D.T. Due, 1986. Myxedematous Cretinism: An Indicator of the Most Severe Goiter Endemias. In: *Frontiers in Thyroidology*, Medeiros-Neto, E. and E. Gaitan (Eds.), Plenum Press, New York, USA., pp: 1081-1084.
- Thomson, B.M., R.W. Vannoort, R.M. Haslemore, 2008. Dietary exposure and trends of exposure to nutrient elements iodine, iron, selenium and sodium from the 2003-4 New Zealand total diet survey. *Br. J. Nutr.*, 99: 614-625.
- Tiwari, B.D., M.M. Godbole, N. Chattopadhyay, A. Mandal and A. Mithal, 1996. Learning disabilities and poor motivation to achieve due to prolonged iodine deficiency. *Am. J. Clin. Nutr.*, 63: 782-786.
- Tonglet, R., P. Bourdoux, T. Minga and A.M. Ermans, 1992. Efficacy of low oral doses of iodized oil in the control of iodine deficiency in Zaire. *N. Engl. J. Med.*, 326: 236-241.
- Van den Briel, T., C.E. West, N. Bleichrodt, F.J. van der Vijver, E.A. Ategbo and J.G. Hautvast, 2000. Improved iodine status is associated with improved mental performance of schoolchildren in Benin. *Am. J. Clin. Nutr.*, 72: 1179-1185.
- Vanderpump, M.P., W.M.G. Tunbridge, J.M. French, D. Appleton and D. Bates *et al.*, 1995. The incidence of thyroid disorders in the community: A twenty-year follow-up of the Whickham survey. *Clin. Endocrinol.*, 43: 55-56.

- Vermiglio, F., M. Sidoti, M.D. Finocchiaro, S. Battiato, V.P.L. Presti, S. Benvenega and F. Trimarchi, 1990. Defective neuromotor and cognitive ability in iodine-deficient schoolchildren of an endemic goiter region in Sicily. *J. Clin. Endocrinol. Metab.*, 70: 379-384.
- Vern, L.C., D.T. Ort and J.L. Grimes, 2003. Physiological factors associated with weak neonatal poulters (*Meleagris gallopavo*). *Int. J. Poult. Sci.*, 2: 7-14.
- Vulsma, T., M.H. Gons and J.J. de Vijlder, 1989. Maternal-fetal transfer of thyroxine in congenital hypothyroidism due to a total organification defect or thyroid dysgenesis. *N. Eng. J. Med.*, 321: 13-16.
- Woods, R.J., A.K. Sinha and R.P. Ekins, 1984. Uptake and metabolism of thyroid hormones by the rat fetus early in pregnancy. *Clin. Sci. (Lond.)*, 67: 359-363.
- Yang, F., Z. Shan, X. Teng, Y. Li and H. Guan *et al.*, 2007. Chronic iodine excess does not increase the incidence of hyperthyroidism: A prospective community-based epidemiological survey in China. *Eur. J. Endocrinol.*, 156: 403-408.
- Zarei, M.A., A. Farshad and S. Akhondzadeh, 2009. Variation in thyroidal activity during estrous cycle and natural breeding season in markhos goat breeds. *Pak. J. Biol. Sci.*, 12: 1420-1424.
- Zhong, F.G., X.M. Cao and J.L. Liu, 1983. Experimental study on influence of iodine deficiency on fetal brain in rats. *Chin. J. Pathol.*, 12: 205-216.
- Zimmermann, M.B., K. Connolly, M. Bozo, J. Bridson, F. Rohner and L. Grimci, 2006. Iodine supplementation improves cognition in iodine-deficient schoolchildren in Albania: A randomized, controlled, double-blind study. *Am. J. Clin. Nutr.*, 83: 108-114.
- Zimmermann, M.B., 2007a. The adverse effects of mild-to-moderate iodine deficiency during pregnancy and childhood: A review. *Thyroid*, 17: 829-835.
- Zimmermann, M.B., 2007b. The impact of iodised salt or iodine supplements on iodine status during pregnancy, lactation and infancy. *Public Health Nutr.*, 10: 1584-1595.
- Zimmermann, M.B., P.L. Jooste, N.S. Mabapa, X. Mbhenyane and S. Schoeman *et al.*, 2007. Treatment of iodine deficiency in school-age children increases insulin-like growth factor (IGF)-I and IGF binding protein-3 concentrations and improves somatic growth. *J. Clin. Endocrinol. Metab.*, 92: 437-442.
- Zimmermann, M.B., 2009. Iodine deficiency in pregnancy and the effects of maternal iodine supplementation on the offspring: A review. *Am. J. Clin. Nutr.*, 89: 668S-672S.
- Zimmermann, M.B. and M. Andersson, 2011. Prevalence of iodine deficiency in Europe in 2010. *Ann. Endocrinol. (Paris)*, 72: 164-166.
- Zou, L., S.C. Hagen, K.A. Strait and J.H. Oppenheimer, 1994. Identification of thyroid hormone response elements in rodent Pcp-2, a developmentally regulated gene of cerebellar Purkinje cells. *J. Biol. Chem.*, 269: 13346-13352.