

Gold Nanoparticle and Mean Inactivation Dose of Human Intestinal Colon Cancer HT-29 Cells

Zahra Arab-Bafrani,^{1,2} Alihossein Saberi,^{3,*} Mohammad Javad Tahmasebi Birgani,² Daryoush Shahbazi-Gahrouei,⁴ Mahdi Abbasian,⁵ and Mehrafarin Fesharaki⁶

¹Disorder Metabolic Research Center, Faculty of Medicine, Golestan University of Medical Sciences, Gorgan, IR Iran

²Department of Medical Physics, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran

³Department of Medical Genetics, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran

⁴Department of Medical Physics and Medical Engineering, School of Medicine, Isfahan University of Medical Sciences, Isfahan, IR Iran

⁵Department of Biotechnology, College of Agriculture, Isfahan University of Technology, Isfahan, IR Iran

⁶Department of Cell Sciences Research Center Medical Science, School of Medicine, Isfahan University of Medical Sciences, Isfahan, IR Iran

*Corresponding author: Alihossein Saberi, Department of Medical Genetics, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran. Tel: +98-9161133488, Fax: +98-613332036, E-mail: saberi-a@ajums.ac.ir; ahsaberi70@hotmail.com

Received 2015 April 16; Revised 2015 June 11; Accepted 2015 June 17.

Abstract

Background: Mean inactivation dose is a useful radiobiological parameter for the comparison of human cell survival curves.

Objectives: Given the importance and accuracy of these parameters, in the present study, the radio sensitivity enhancement of colon cancer (HT-29) cells in the presence of gold nanoparticles (GNPs) were studied using the mean inactivation dose (MID).

Materials and Methods: Naked-GNPs with 50 nm diameters were incubated with HT-29 cells. The cytotoxicity and uptake of these particles on HT-29 cells were assessed. After determining the optimum GNPs concentration, the cells were incubated with gold nanoparticle for 24 hours. The change in the MID value as well as the radio sensitization enhancement under irradiation with 9 MV X-ray beams in the presence of GNPs were evaluated by multiple (3-(4, 5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium) MTS assay.

Results: Cell survival in the presence of GNPs was more than 90% and the maximum uptake of GNPs was observed at 60 μ M of gold nanoparticles. In contrast, in the presence of GNPs combined with radiation, cell survival and MID value significantly decreased, so that the radio sensitization enhancement was 1.4.

Conclusions: Due to the significant reduction in the mean inactivation dose of colon cancer cells in the presence of gold nanoparticles, it seems that GNPs are suitable options to achieve a new approach in order to improve radiotherapy efficiency without increasing the prescribed radiation dose.

Keywords: Colon Cancer, Gold Nanoparticle, Mean Inactivation Dose, HT-29 Cells

1. Background

Colorectal cancer is the second leading cause of cancer-related mortalities in developed countries. In Asian and middle eastern countries, due to changes in dietary habit and lifestyle, the incidence of colon cancer has shown a remarkable increase. Nowadays, surgery and chemotherapy are the top choices for colorectal cancer treatment and radiation therapy is the complementary therapy (1-3). However, despite all the recent developments in cancer therapy, colon cancer recurrence (50%) has remained as a major problem and it is necessary to develop some new more effective approaches for achieving a certain cell mortality rate (3, 4). Increasing the irradiation dose improves the efficiency of treatment, but the side effects of high-dose radiation including the damage of normal tissues are among the limitative factors (5). The application of radio sensitizers is one of the strategies to enhance the radiation efficiency without exceeding the maximum tolerable

dose of normal tissues. Today, gold nanoparticles (GNPs) are established as good radio sensitizers due to their high bio-compatibility and high penetration rate into cancer cells (6, 7). In addition, GNPs can be conjugated with antibodies or other proteins to target tumor cells as well as the intracellular targets such as the nucleus (8, 9). In fact, the presence of GNPs in radiation events results in local dose amplification and radiosensitization enhancement. The interactions of radiation elements with these particles lead to the production of more secondary electrons. These electrons include photo and Compton electrons with high energies. Followed by these events, a shower of Auger electrons with low energy and small range are produced. Small range of Auger electrons let them deposit their energy in the vicinity of GNPs, which leads to increased cell damage and amplified local doses (6, 10-13).

The relationship between radiation dose and radiobio-

logical parameters can be derived with cell survival data. Linear quadratic (LQ) model is widely used to analyze cell survival data and is useful to investigate the radio resistance mechanisms and design new therapeutic trends. With this model, the radiobiologic parameters that describe the behavior of cells over the whole survival curve are accessible and also significant differences in radio-sensitivity that exist among histological categories of human cell lines can be determined.

Mean inactivation dose (MID), one of radiobiological parameters, is a useful concept for the comparison of human cell survival curve. Unlike other parameters of radiobiology, MID estimates the radiation sensitivity of mammalian cells at low and high radiation doses (14). Less dependence to deviation of the survival curve of a specific cell line which was investigated by different researchers and usability for cells with different histological categories are the advantages of MID rather than other radiobiological components such as the multi-target parameters D_0 and n . Accordingly, international commission on radiation units (ICRU) recommend the use of MID to characterize the cell survival curve (14, 15).

2. Objectives

Given the importance of discovering an effective method to cure colon cancer, the emergence of new models that suggest radiation dose enhancement in the presence of GNPs, and the advantages of MID as an indicative parameter of changes in cell radiosensitivity, in this study, the effect of gold nanoparticle on MID value of colon cancer cells at X-ray irradiation was investigated.

3. Materials and Methods

3.1. Cell Culture

Colorectal (HT-29) cell line was purchased from Pasteur institute (Tehran, Iran). Cells were cultured in Roswell park memorial institute (RPMI) 1640 medium (Gibco-Invitrogen), supplied by 10% fetal bovine serum (Gibco-Invitrogen), 1% antibiotic mixture containing penicillin (Sigma-Aldrich), and streptomycin (Sigma-Aldrich). The cells were stored at humidified atmosphere at 37°C with 5% CO₂. The medium of cells was changed approximately every two days, and when they reached more than 80% confluency, they were split with 0.05% trypsin/0.02% EDTA and sub-cultured for more passages.

3.2. Gold Nanoparticles Cytotoxicity and MTS Assay

The cells were seeded at 2×10^3 per well of a 96-well tissue culture plate. After 48 hours, GNPs (10 and 100 μM) were added to the cells. Cell viability was investigated using MTS assay, the CellTiter 96 AQueous One Solution Cell Proliferation (ProMega), after 2, 4, 6, 24 and 48 hours. The absorbance was recorded at 490 nm in a 96-well plate reader (Biorad). Cell survival was evaluated using the following equation (Equation 1) (16).

$$(1) \text{ Survival Fraction} = \frac{\text{mean OD in test wells} - \text{mean OD in cell free wells}}{\text{mean OD in control wells} - \text{mean OD in cell free wells}}$$

3.3. Uptake Assay of Gold Nanoparticle

When cells reached more than 80% confluency, they were trypsinized and seeded in a 24-well plate (4000 cells/well) for 48 hours (in logarithmic phase). Consequently, the cells were exposed to 20, 40, 60, 80 and 100 μM of a 50 nm-gold nanoparticle (Sigma-Aldrich) and stored at humidified atmosphere at 37°C with 5% CO₂ for 24 hours. When the incubation period passed, the culture medium was removed, the plate was washed three times with phosphate-buffered saline (PBS), the cells were counted, and 1 mL HCL (5 M) was added to each well for lysis of the cells. The concentration of GNPs was measured by graphite furnace atomic absorption spectrometry (GFAAS) analysis.

3.4. Irradiation

A total of 2×10^3 cells per well were plated in 96-well plates for 48 hours. Then, they were incubated with the gold nanoparticle for 24 hours before irradiation. Irradiation was performed using a clinical accelerator (Neptun 10 PC) at a source-to-surface distance (SSD) = 100 cm and 20×20 cm² field size. Two centimeters of a Plexiglass sheet (water equivalent) was placed on the top of the plate to serve as a built-up material for the 9 MV beam. A Plexiglass sheet (water equivalent) of 5 cm was placed under the bottom of the plate to ensure the full backscatter condition (Figure 1). Mega-voltage radiation (9 MV) was delivered at total doses of 2, 4, 6 and 8 Gy with a dose rate of 300 cGy min⁻¹. In vivo radiation diode dosimetry measurements were performed for beam calibration and variation within a field was smaller than 2% for each well. After irradiation, the cells were incubated at 37°C with 5% CO₂. All the treatments were carried out in triplicate and were repeated.




Figure 1. Schematic Cross-Section of the Cell Irradiation Phantom

3.5. Survival Fraction-Based Multiple-MTS Assay

Multiple-MTS assay was used to estimate the survival fraction (17). In multiple-MTS assay, the cell growth data (the number of viable cells) are collected daily to draw the cell growth curve. The survival fraction can be estimated by measuring the time difference between the cell growth of the control and the treated groups to reach a certain value. Using the CellTiter 96 AQueous One Solution Cell Proliferation Assay (ProMega), cell viability tests were performed directly after irradiation (1 - 7 days) and cell growth curve was obtained.

Survival fraction was calculated by the following equation (Equation 2) (17):

$$(2) \quad \text{Survival} = 2^{-\frac{t_{\text{delay}}}{t_{\text{doubling time}}}}$$

Where t doubling time is the period required for a quantity of cells to double and t delay is the period to reach the specific absorption value of control versus irradiated cells.

3.6. Evaluation of Mean Inactivation Dose and Radio Sensitization Enhancement in the Presence of Gold Nanoparticle

MID is equal to the area under the cell survival curve. After estimating the survival fraction at different radiation doses, the survival curve (survival fraction versus radiation dose) should be fitted to the linear quadric model (Equation 3). MID was calculated using the following equation (Equation 4) (14).

$$(3) \quad S = e^{-\alpha D - \beta D^2}$$

$$(4) \quad \text{MID} = \int S(D) d(D)$$

The Sensitizer enhancement ratio (SER) value was calculated by dividing the MID of cells not exposed to GNPs to cells exposed to GNPs.

4. Results


4.1. Gold Nanoparticles Uptake in HT-29 Cells

HT-29 was incubated with GNPs in varying concentration (20, 40, 60, 80 and 100 μM). Figure 2 shows the average gold nanoparticle uptake by each cell in an individual concentration. It indicates that the uptake of GNPs increases with increase in GNPs concentration. The maximum uptake was observed in 80 μM (62.98 ± 3.49) $\times 10^3$ and there was no significant difference in GNPs uptake between 80 μM and 100 μM (P-value = 0.1 > 0.05).

4.2. Cytotoxicity of Gold Nanoparticles


To determine the cytotoxicity of GNPs on HT-29 cells, the cells were incubated with different concentrations of GNPs (10, 20, 50 and 100 μM) for 24 hours. The cytotoxicity of GNPs at different concentrations was measured by MTS assay. Figure 3 a shows no significant difference between the controls and those treated with the GNPs over the whole range of GNP concentrations (P value > 0.05). In addition, because the cytotoxicity of nanoparticles could be time-dependent, MTS assay was performed at different incubation times (2, 4, 6, 24 and 48 hours). Cell viability at different times was over 90% in all the groups (P > 0.05) (Figure 3 b). Collectively, this data indicated that GNPs did not have remarkable cytotoxicity on HT-29 cells in all the studied concentrations (10 - 100 μM).

Figure 2. Study of Cellular Uptake of Gold Nanoparticles at Different Concentration After Incubating with GNPs for 24 Hours




Cellular uptake of gold nanoparticles is as a function of concentration.


Figure 3. Survival Fraction of HT-29 Cells, (left) Incubated With Different GNP Concentrations (10, 20, 50 and 100 μM) at a Fixed Incubation Time; (right) at Various Time Periods With 10 and 100 μM Concentration



Cell viability in all the groups at different times was over 90% (P > 0.05).

Figure 4. The Effect of Gold Nanoparticles During Irradiation on the Growth of Irradiated Cells at 7 Days After X-Ray Irradiation

A, alone irradiation; B, GNPs irradiation.

Figure 5. The Survival Curves of HT-29 Cells in the Presence and Absence of Gold Nanoparticles Irradiated With 9 MV X-Ray Beam

Solid line, alone-irradiation; dash line, GNPs irradiation.

4.3. Cell Survival Fraction Curve in the Presence of Gold Nanoparticles

For evaluating the effectiveness of GNPs on radiosensitivity of HT-29 cells, they were incubated with a 60 μ M concentration of GNPs for 24 hours and then were exposed to 2, 4, 6 and 8 Gy of 9 MV X-ray energy. Figure 4 Shows that GNP-irradiated cells decreased more than alone-irradiated cells. Multiple-MTS assay was used to quantify the survival fraction of irradiated cells in the presence and absence of GNPs. MTS assay was conducted in several days and cell growth curve was drawn for control, GNPs alone, X-ray alone, and X-ray with GNPs groups. The survival fraction of alone irradiated cells and GNPs-irradiated cells was estimated based on the delay that occurred between growth curve of the control and treated cells.

Figure 5 shows the survival curve of the two mentioned groups based on the LQ-model. As can be noted from Figure 5, radiation combined with GNPs induced greater decreases in survival fraction at different radiation doses ($P < 0.05$).

4.4. Effect of Gold Nanoparticles on Mean Inactivation Dose

After drawing the cell survival curve, MID value was estimated using Equation 4. The MID value in cells only irradiated was 5.03 and significantly reduced in cells irradiated in the presence of GNPs (3.61).

Sensitive enhancement ratio (SER) that was calculated by dividing the MID value to the two mentioned groups, was 1.4.

5. Discussion

Cell radiosensitivity can be predicted by radiobiological parameters extracted from the survival curve. Mean inactivation dose is one of these parameters that is introduced for characterizing of cell radiosensitivity over the whole survival curve with a single parameter. The usability for the whole cell population and the presentation of the consistency results make this parameter more acceptable for describing cell radio sensitivity compared to other sets of radiobiological parameters (14, 18, 19).

In several theoretical and experimental studies, using GNPs as radiosensitizers at orthovoltage (kV) and megavoltage (MV) energies has been demonstrated by amplifying local radiation dose and consequently increasing the rate of cell death (20-22). Recently, researchers reported that GNPs can enhance the radiosensitization of human carcinoma cells such as prostate, ovarian and breast cancer cells. Jain et al. (10) demonstrated that GNPs radiosensitization depended on the type of cell. Our results indicated that GNPs irradiation induced greater decreases

in survival fraction than alone irradiation at different radiation doses and enhanced the radiosensitization of colon cancer cells. Based on previous studies and given the importance of MID parameter to predict radiosensitization, the present study aimed to calculate the SER of colon cancer cells by dividing MID of cells not exposed to GNPs to cells exposed to GNPs.

The cytotoxicity and the uptake of GNPs are two important parameters that influence radiosensitization (12, 23). Our results clearly revealed that cell viability was more than 90% when colon cancer cell (HT-29) was incubated with GNPs at 10 - 100 μ M. Based on the steps of the uptake assay, GNPs on the cell surface were removed after three times washing with PBS and the average numbers of naked-GNPs uptake in 60 μ M and 80 μ M were $(5.749 \pm 5.65) \times 10^4$ and $(6.298 \pm 3.49) \times 10^4$, respectively, and there was no significant difference in uptake ($P = 0.09 > 0.05$). Wang et al. (24) showed that the amount of Glu-GNPs in A549 cells was approximately 12×10^4 and demonstrated that significant radiosensitization occurred in A549 cells with an SER of 1.49 at 6 MV X-rays with Glu-GNPs. In another study on MDA-MB-231 cells (10), the SER was found to be 1.29 and 1.16 for 6 MV and 15 MV irradiations, respectively, and in a similar study (25), the corresponding values were 1.24 ± 0.05 and 1.18 ± 0.04 . Taken together these data, determining SER-based MID value is a useful approach for predicting the radiosensitizing effects of GNPs. In agreement with previous studies, the obtained results revealed a significant reduction in survival fraction of HT-29 cells that were exposed to 9 MV X-ray with GNPs over the whole survival curve ($P < 0.05$). The MID value of HT-29 cells that were incubated with GNPs (3.61) significantly decreased rather than the ones not exposed to GNPs (5.032) and the SER was 1.41.

In conclusion, our results suggested that in the presence of GNPs, the MID value of HT-29 cells reduced and consequently GNPs can induce radiosensitization enhancement in colon cancer cells.

Acknowledgments

This study was the result of a PhD project carried out by the first author under the supervision of the second author and with the help and advice of other co-authors as the advisers at Jundishapur university of medical sciences, Ahvaz, Iran. Hence, we would like to express our special thanks to these institutions for providing us the financial, technical and clinical support required for carrying out this research.

Footnotes

Authors' Contribution: Alihossein Saberi, Zahra Arab-Bafrani and Mohammad Javad Tahmasemi Birgani developed the original idea and the protocol, abstracted and analyzed data, wrote the manuscript, and were the guarantors. Other authors contributed to the development of the protocol, abstracted the data and prepared the manuscript.

Funding/Support: This work was financially supported by Ahvaz Jundishapur university of medical sciences, Ahvaz, Iran.

References

1. Azadeh S, Moghimi-Dehkordi B, Fatem SR, Pourhoseingholi MA, Ghiasi S, Zali MR. Colorectal cancer in Iran: an epidemiological study. *Asia Pac J Cancer Prevent*. 2007;9(1):123-6.
2. Albuquerque C. The 'just-right' signaling model: APC somatic mutations are selected based on a specific level of activation of the beta-catenin signaling cascade. *Human Mol Gene*. 2002;11(13):1549-60. doi: 10.1093/hmg/11.13.1549.
3. Chen WS, Lee YJ, Yu YC, Hsiao CH, Yen JH, Yu SH, et al. Enhancement of p53-mutant human colorectal cancer cells radiosensitivity by flavonoid fisetin. *Int J Radiat Oncol Biol Phys*. 2010;77(5):1527-35. doi: 10.1016/j.ijrobp.2010.02.043. [PubMed: 20637980]
4. Xu Y. Induction of genetic instability by gain-of-function p53 cancer mutants. *Oncogene*. 2008;27(25):3501-7. doi: 10.1038/sj.onc.1211023. [PubMed: 18223686]
5. Andreyev HJ. Gastrointestinal problems after pelvic radiotherapy: the past, the present and the future. *Clin Oncol (R Coll Radiol)*. 2007;19(10):790-9. doi: 10.1016/j.clon.2007.08.011. [PubMed: 17904338]
6. Jain S, Hirst DG, O'Sullivan JM. Gold nanoparticles as novel agents for cancer therapy. *Br J Radiol*. 2012;85(1010):101-13. doi: 10.1259/bjr/59448833. [PubMed: 22010024]
7. Smith L, Kuncic Z, Ostrikov K, Kumar S. Nanoparticles in Cancer Imaging and Therapy. *J Nanomater*. 2012;2012:1-7. doi: 10.1155/2012/891318.
8. Jiang W, Kim BY, Rutka JT, Chan WC. Nanoparticle-mediated cellular response is size-dependent. *Nat Nanotechnol*. 2008;3(3):145-50. doi: 10.1038/nnano.2008.30. [PubMed: 18654486]
9. Nativo P, Prior IA, Brust M. Uptake and intracellular fate of surface-modified gold nanoparticles. *ACS Nano*. 2008;2(8):1639-44. doi: 10.1021/nn800330a. [PubMed: 19206367]
10. Jain S, Coulter JA, Hounsell AR, Butterworth KT, McMahon SJ, Hyland WB, et al. Cell-specific radiosensitization by gold nanoparticles at megavoltage radiation energies. *Int J Radiat Oncol Biol Phys*. 2011;79(2):531-9. doi: 10.1016/j.ijrobp.2010.08.044. [PubMed: 21095075]
11. McMahon SJ, Hyland WB, Muir MF, Coulter JA, Jain S, Butterworth KT, et al. Biological consequences of nanoscale energy deposition near irradiated heavy atom nanoparticles. *Sci Rep*. 2011;1:18. doi: 10.1038/srep00018. [PubMed: 22355537]
12. Rahman WN, Bishara N, Ackerly T, He CF, Jackson P, Wong C, et al. Enhancement of radiation effects by gold nanoparticles for superficial radiation therapy. *Nanomed Nanotechnol Biol Med*. 2009;5(2):136-42. doi: 10.1016/j.nano.2009.01.014.
13. Xiao F, Zheng Y, Cloutier P, He Y, Hunting D, Sanche L. On the role of low-energy electrons in the radiosensitization of DNA by gold nanoparticles. *Nanotechnology*. 2011;22(46):465101. doi: 10.1088/0957-4484/22/46/465101. [PubMed: 22024607]
14. Fertil B, Dertinger H, Courdi A, Malaise EP. Mean inactivation dose: a useful concept for intercomparison of human cell survival curves. *Radiat Res*. 1984;99(1):73-84. [PubMed: 6739728]
15. Belli M, Simula S. The calculation of the mean inactivation dose by approximated methods. *Radiat Res*. 1991;128(1):112-4. [PubMed: 1924722]
16. Price P, McMillan TJ. Use of the tetrazolium assay in measuring the response of human tumor cells to ionizing radiation. *Cancer Res*. 1990;50(5):1392-6. [PubMed: 2302704]
17. Buch K, Peters T, Nawroth T, Sanger M, Schmidberger H, Langguth P. Determination of cell survival after irradiation via clonogenic assay versus multiple MTT Assay—a comparative study. *Radiat Oncol*. 2012;7:1. doi: 10.1186/1748-717X-7-1. [PubMed: 22214341]
18. Brenner DJ. The linear-quadratic model is an appropriate methodology for determining isoeffective doses at large doses per fraction. *Semin Radiat Oncol*. 2008;18(4):234-9. doi: 10.1016/j.semradonc.2008.04.004. [PubMed: 18725109]
19. Weichselbaum RR, Beckett MA, Vijayakumar S, Simon MA, Awan AM, Nachman J, et al. Radiobiological characterization of head

- and neck and sarcoma cells derived from patients prior to radiotherapy. *Int J Radiat Oncol Biol Phys.* 1990;**19**(2):313-9. doi: 10.1016/0360-3016(90)90539-v. [PubMed: 2394610]
20. Cho SH. Estimation of tumour dose enhancement due to gold nanoparticles during typical radiation treatments: a preliminary Monte Carlo study. *Phys Med Biol.* 2005;**50**(15):N163-73. doi: 10.1088/0031-9155/50/15/N01. [PubMed: 16030374]
21. Roa W, Zhang X, Guo L, Shaw A, Hu X, Xiong Y, et al. Gold nanoparticle sensitize radiotherapy of prostate cancer cells by regulation of the cell cycle. *Nanotechnology.* 2009;**20**(37):375101. doi: 10.1088/0957-4484/20/37/375101. [PubMed: 19706948]
22. Kong T, Zeng J, Wang X, Yang X, Yang J, McQuarrie S, et al. Enhancement of radiation cytotoxicity in breast-cancer cells by localized attachment of gold nanoparticles. *Small.* 2008;**4**(9):1537-43. doi: 10.1002/sml.200700794. [PubMed: 18712753]
23. Patra HK, Banerjee S, Chaudhuri U, Lahiri P, Dasgupta AK. Cell selective response to gold nanoparticles. *Nanomedicine.* 2007;**3**(2):111-9. doi: 10.1016/j.nano.2007.03.005. [PubMed: 17572353]
24. Wang C, Li X, Wang Y, Liu Z, Fu L, Hu L. Enhancement of radiation effect and increase of apoptosis in lung cancer cells by thio-glucose-bound gold nanoparticles at megavoltage radiation energies. *J Nanopart Res.* 2013;**15**(5):1-12. doi: 10.1007/s11051-013-1642-1.
25. McMahon SJ, Hyland WB, Muir MF, Coulter JA, Jain S, Butterworth KT, et al. Nanodosimetric effects of gold nanoparticles in megavoltage radiation therapy. *Radiother Oncol.* 2011;**100**(3):412-6. doi: 10.1016/j.radonc.2011.08.026. [PubMed: 21924786]